

**Nutritional and toxicological studies on New Zealand
Mutton bird meat (*Puffinus griseus*)**

A thesis submitted in partial fulfillment
of the requirements for the degree
of
Master of Science

at
Lincoln University
by
Saleh Suliaman Al-Amer

Lincoln University

2009

Abstract & Keywords:

An abstracts of a thesis submitted in partial fulfillment of the requirements for the degree of
Master of Science (M. Sc)

Nutritional and toxicological studies on New Zealand

Mutton bird meat (*Puffinus griseus*)

by

Saleh Al-Amer

New Zealand mutton bird or tītī (*Puffinus griseus* and order procellariiformes) nest in New Zealand during the summer months, migrate to the northern hemisphere during May and return in September. Their eggs are laid during November and December and the chicks are hatched in the following January and February. Large numbers of them are harvested from April to May in New Zealand. They are wild seabirds annually harvested by Maori according to the customary rights agreement set by Treaty of Waitangi.

NZ mutton birds also called Sooty Shearwaters are noted for their high proportion of body fat. These birds are interesting since its sole diet is based on krill and other small marine organisms that are potentially rich in n-3 fatty acids and other marine bioactive compounds. The proximate composition, fatty and amino acids and cholesterol content of mutton bird pectoral muscle were determined and compared with other common meat to explore the nutritional value of this New Zealand delicacy. The concentration of twenty two essential and toxic elements including silver (Ag), aluminium (Al), arsenic (As), cadmium (Cd), chromium (Cr), copper (Cu), mercury (Hg), manganese (Mn), lead (Pb), selenium (Se), potassium (K), calcium (Ca), magnesium (Mg), boron (B), iron (Fe), nickel (Ni), sulphur (S), sodium (Na) and cobalt (Co) and zinc (Zn) in mutton bird breast meat (*Puffinus griseus*) were measured over two years to evaluate its safety for human consumption. Persistent organochlorine, dichlorodiphenyltrichloroethane (DDT) and their metabolites, and aldrin and lindane were also measured.

Twenty bird carcasses were purchased in both 2006 and 2007 from a local source. Meat samples from the pectoral muscle of two carcasses were pooled to generate 10 samples for each year. These were used for trace element analysis using inductively coupled plasma-mass spectrometry (ICP-MS). Trace elements were in the range of 0 to 1.09 mg/kg wet weight for Ag, 0 to 3.32 for Al, 0.17 to 0.79 for As, 0.01 to 0.07 for Cd, 0.03 to 0.15 for Cr, 3.56 to 4.88

for Cu, 0 to 0.15 for Hg, 0.22 to 0.50 for Mn, 0 to 0.09 for Pb, 0.66 to 1.18 for Se and 11.49 to 23.70 for Zn. In 2006, Ag, Al, Mn and Zn concentrations were significantly higher but Pb and Hg concentrations significantly lower compared to the 2007 samples ($P < 0.05$). Apart from one sample in 2006, all the samples were below the published maximum level for concern. However, our preliminary data indicated that the higher level of Cd and other metals in the skin of mutton bird may compromise the overall safety to humans consuming the skin of mutton birds. It is suggested that the evaluation of the metals in different parts and/or the whole mutton bird at different seasons is required to assure complete safety to the consumers.

Furthermore, the nutritional value of mutton bird meat was studied over two harvesting seasons (2006 and 2007) to investigate the impact of seasonal variation. The moisture and carbohydrates contents ranged between 54.0 to 55.0 % and 2.8 to 3.0 %, respectively, and no seasonal effects were evident in these components. The values for fat and ash contents were higher and the protein content lower for birds harvested in 2007 compared with the 2006 values which ranged from 11.8 to 13.0, 10.3 to 11.7, and 20.3 to 18.5 % for fat, ash and protein content respectively. The major amino acids in mutton bird pectoral muscle were glutamate, aspartate, lysine, leucine, and arginine. Higher lysine concentrations and lower proline, cysteine and methionine were found in mutton birds compared with the literature values for beef, lamb and pork. The essential amino acid content in mutton bird (43.8 and 44.9 % in 2006 and 2007, respectively) was slightly higher than those found in beef and lamb meats (42-43%).

The major fatty acids detected were palmitic (C16:0), stearic (C18:0), oleic and isomers (C18:1), eicosenoic (C20:1), Docosahexaenoic acid (DHA) (C22:6), icosapentaenoic acid (EPA) (C 20:5) and these accounted for approximately 77% of the fatty acids. The $\omega 3/\omega 6$ ratio of fats from pectoral muscle was 1.3. The cholesterol concentration varied slightly in the two years with 184.4 ± 37.37 and 134.4 ± 25.55 mg/100 g fresh weight for 2007 and 2008 respectively. Mutton bird was shown to contain significantly higher cholesterol content (134.4-184.4) than other common meat such as chicken (80.3-88.9), lamb (62.3), fish (52.79) and beef (51.97). Overall, the nutritional value of mutton bird muscle was similar to or superior to the traditionally protein sources such as seafood and red meat. Annual variations existed in the composition of Mutton bird pectoral muscle but this is not of nutritional consequence but might be a useful indicator for ecological events such as feed availability and other environmental issues. Mutton bird seems to be a good source of essential minerals, Zn and Fe compared with other traditional meats source. Mutton bird meat is nutritionally as good as the major sources of red or white meats. It may even have advantages over the other common meats (beef, lamb, fish and chicken) due to its high protein and monounsaturated fatty acids (omega n-3 and n-6) content. However, its high cholesterol content may represent a risk factor for some people.

Keywords: mutton bird, trace elements, minerals, fatty acids, amino acids, cholesterol, seasonal variation breast meat, nutrition and toxicology.

Acknowledgements

Definitely, this thesis as any study was made and supported by kind help from many people to whom I am really grateful.

Firstly, I would like to thank very much my supervisor, associate. Professor .Ravi Gooneratne who was patient during my study and he was continuously encouraging, helpful and faithful. Deeply I am indebted to him for his ongoing assistance and direction throughout the duration of this work and spending a lot of his valuable time in order to read, organize and correct my drafts. To my associate supervisor, Dr Sue Mason also I am very thankful for her good advice and teaching particularly the nutritional aspect of my study as she provided me with support in the experimental work and assisted in proof reading and correcting all drafts.

Indeed, I extend my personal thanks to my teacher preferred Dr. Alaa- El-din A Bekhit who is my external supervisor. His immense encouragement and support during my studies and research is gratefully acknowledged. His encouragement and explanation when I am having difficulties lifted me up and made me confident. I also like to thank both Dr. Jim Morton and Dr. Mike Noonan for their good teaching during my course work. I would like to express my profound gratitude to all technicians in our department who trained me and were kindly helpful throughout my research, in particular Jenny Zhao, Chris Dawson, Jeff Thackwell, Martin Wellby and Lynne Clucas. I would also like to express my gratitude to Dr. Khalid Othman, Al-Qassim university, Saudi Arabia for giving me space in his laboratory to carry out the DDT analysis during my holidays at Al-Qassim from December 2008 to February 2009.

I take this opportunity to thank my classmate, Mathew Jicheha for his support throughout the duration of my research. A special thank to my fellow student, Jinlin Shi for her help during my research. Finally, I would like to convey my warm thanks to the administrative secretaries, particularly Marianne Duncan and Jan Haldane for their lovely reception, help and sense of humors.

Contents	Page
Declaration	II
Abstract	III
Acknowledgments	V
Contents	VI
List of Tables	X
List of Figures	XII
Chapter 1 Introduction	1
Chapter 2 Literature Review	4
2.1. Overview of Mutton bird	
2.1.1 Procellariidae family	
2.1.2 Sooty shearwater	
2.1.3 Location and distribution of Mutton bird in New Zealand	6
2.1.4 Characteristics and life cycle of Mutton bird	7
2.1.5 The industry and Maori methods	8
2.1.6 The relationships between Mutton-birds and krill	9
2.2 Nutrition of Avian meat	10
2.2.1 Mutton bird	
2.2.2 Chicken meat quality	13
2.2.2.1 Proximate composition of chicken meat	14
2.2.2.2 Fatty acids in chicken	15
2.2.2.3 Nutritional benefits of fatty acids	16

2.2.2.4	The relationship between fatty acids and cholesterol with lipid oxidation	17
2.2.2.5	The correlation between fatty acids and cholesterol	18
2.2.2.6	The role of Vitamin E in meat quality	
2.2.2.7	Amino acids in chicken meat	20
2.3	Toxicological aspects	21
2.3.1	Trace elements in sea birds	
2.3.2	The use of seabirds as biomonitors for environmental pollution	22
2.3.3	Pesticides in sea birds	25
2.3.4	Distribution and occurrence of toxic compounds in seabirds.	27
2.3.4.1	Heavy metals	
2.3.4.2	Organochlorine	30
2.3.4.3	Residual organochlorine pesticides in Mutton bird.	31
2.4	Heavy metals in chicken	32
2.5	Hypotheses and objectives	33
2.5.1.	Hypothesis	
2.5.2	Objectives	
2.5.2.1	Broad Objectives	
2.5.2.2	Specific Objectives	
Chapter 3	Nutritional studies on Mutton bird meat (<i>Puffinus griseus</i>)	34
3.1	Introduction	
3.2	Material and Methods	38

3.2.1	Sample collection	
3.2.2	Size and weight measurements:	
3.2.3	Sample preparation for Analysis:	
3.2.4	Proximate composition	39
3.2.4.1	Moisture analysis	
3.2.4.2	Crude fat analysis	
3.2.4.3	Crude protein (total nitrogen) analysis	
3.2.4.4	Ash analysis	40
3.2.4.5	Carbohydrate analysis	
3.2.5	Fatty acid analysis	
3.2.5.1	Extraction and preparation of samples	
3.2.5.2	Extraction of oil	
3.2.5.3	Preparation of Methyl esters for Fatty acids analysis.	41
3.2.5.4	Fatty acid separation and detection	42
3.2.5.5	Standards and quantification	43
3.2.6	Amino acid analysis	
3.2.6.1	Preparation and extraction of samples	
3.2.6.2	HPLC analysis of amino acid	44
3.2.6.3	Standards and quantification	46
3.2.7	Cholesterol analysis	47
3.2.7.1	Method of cholesterol analysis	
3.2.7.2	Preparation of samples and standards	
3.2.7.3	Cholesterol analysis	

3.2.8 Statistical analysis	48
3.3 Results and discussion	
3.3.1 Proximate composition	
3.3.2 Amino acid composition of Mutton bird breast meat.	50
3.3.2.1 Amino Acid composition for samples collected in 2007and 2008.	
3.3.3 Fatty acid composition	54
3.3.4 Cholesterol content	61
Chapter 4 Toxicological studies on Mutton bird	64
4.1 Introduction	
4.2 Materials and methods	67
4.2.1 Determination of elements in Mutton bird pectoral muscle.	
4.2.2 Determination of Hg in Mutton bird muscle	68
4. 3 DDTs analysis	70
4.3.1 Results and discussion	74
4.3. 2 Trace elements in Mutton bird breast meat	74
4.3.2.1 Organochlorines in Mutton bird breast meat	90
Chapter 5 General Discussion	95
5.1 Introduction	
5.2 Nutritional values in MBBM	96
5.3 Toxicological studies in MBBM	101
5.4 Conclusion	102
Chapter 6 References	103

List of Tables	Page
Table 2.1: Main characteristics of sooty shearwater (Richdale, 1976).	7
Table 2.2: Fatty Acid Profile of NZ Mutton bird by Quigley et al., 1995.	11
Table 2.3: Composition (per 100 g edible portions) of cooked skinless Mutton bird flesh (Woodward et al., 1995).	12
Table 2.4: Fatty acid profile of skinless Mutton bird flesh lipid (Woodward et al., 1995).	12
Table 2.5: Proximate composition mg/100g and lipids in various parts of chicken (Mueller, 2006).	14
Table 2.6: Amino acid composition of red and white meats (Mavromichalis et al., 2000).	20
Table 2.7: Concentrations mg.kg ⁻¹ of eight trace elements in seven regions in Nigeria (Iwegbue et al., 2008).	32
Table 3.1: Preperation of Standards (1mg/ml) for Liberman-Burchard method of cholesterol estimation (mls).	47
Table 3.2: Mean (standard deviation) of moisture, crude fat, ash, and protein and carbohydrate content of Mutton bird pectoral muscle obtained in 2007 and 2008.	48
Table 3.3: Proximate compositions of different animal meat.	49
Table 3.4: A comparison of amino acid composition for Mutton bird breast between years (2007 and 2008) (g/100g fresh weight).	51
Table 3.5: A comparison of amino acid composition for Mutton bird breast between years (2007 and 2008) (g/ 100 g protein).	52
Table 3.6: A comparison of the amino acid content of Mutton bird breast meat with other common meats (g/100g of protein).	53
Table 3.7: Fatty acid compositions of Mutton birds' breast for two years (2007 & 2008).	55
Table 3.8: Comparison of Mutton bird fatty acid composition breast meat with other common meats.	57
Table 3.9: Mean and standard deviation for cholesterol content of Mutton Bird's breast meat mg/100 fresh weight for years 2007 and 2008.	62

Table 3.10: Comparison of cholesterol content in Mutton bird meat with other common meats.	62
Table 4.1: Retention times, molecular weight and fragment ions selected for confirmation compounds using GC-MS.	73
Table 4.2: Heavy metal concentrations in New Zealand Mutton bird breast meat for 2007 and 2008.	75
Table 4.3: Trace element concentrations of Mutton bird breast meat and other common meats.	77
Table 4.4: Levels (ng/g = ppb) of organochlorines in the Mutton bird breast meat collected in two years (2007-2008).	91
Table 4.5: A comparison of DDT type compounds (DDT+ DDE+ DDA) between Mutton breast bird muscle and other common meats.	92

List of Figures	Page
Figure 2.1: Main islands for the traditional harvest of Mutton bird in the south of New Zealand (Kitson et al., 2000).	5
Figure 2.2: The route of Mutton bird migration to NZ (Anderson, 1996).	7
Figure 2.3: shown the sooty shearwater <i>U.S. Fish & Wildlife Service – Alaska, 2006</i>).	7
Figure 2.4: The food web of the Southern Ocean with the krill being the basal diet for all marine organisms (Everson, 2000).	10
Figure 2.5: shows the abbreviations and chemical configurations of common fatty acids (Bayır et al., (2006).	16
Figure 2.6: Structure of tocopherol and tocotrienol (Kramer et al., 1997).	19
Figure 2.7: Distribution of element burdens in tissues and organs of three seabirds (Kim et al. 1998).	28
Figure 3.1: A representative sample of Mutton bird.	38
Figure 3.2: Proximate analysis of Mutton bird and other common meats.	50
Figure 3.3: Amino acids for Mutton bird and other common white meats.	54
Figure 3.4: Amino acids in Mutton bird breast meat and common red meats.	54
Figure 3.5: The fatty acid contents in different common red and white meats.	61
Figure 3.6: Comparison of cholesterol content of Mutton bird muscle with other common white and red meats.	63
Figure 4.1: Show amounts of Aldrin in Mutton bird muscle and other common meats.	93
Figures 4.2: Show amounts of Lindane in Mutton bird muscle and other common meats.	94