

Can topdressing with salt increase oversowing success and pasture quality on steep, south facing slopes in hill country pastures?

B.J. GILLESPIE, R.J. LUCAS, D.J. MOOT and G.R. EDWARDS

Agriculture Group, Agriculture and Life Sciences Division, Field Service Centre, Lincoln University, Lincoln 7647
edwardsg@lincoln.ac.nz

Abstract

Two experiments were conducted in sodium deficient (<0.03% Na in DM) pastures on steep, south facing slopes at Mt Grand, Hawea, Central Otago (600 m.a.s.l.), to determine the effect of the application of coarse salt (NaCl) fertiliser on pasture composition and establishment of oversown seeds. In Experiment 1, factorial combinations of salt (0, 150 kg NaCl/ha), N (0, 100 kg N/ha as urea) and sulphur superphosphate (0, 500 kg/ha) fertilisers were applied to 30 x 8 m plots in early December 2003. Balansa clover (10 kg seed/ha) and subterranean clover (10 kg/ha) seeds were oversown into each plot in March 2004. Plots where salt was applied were grazed intensively by Merino ewes immediately after salt application creating up to 50% bare ground. In December 2004, pastures where salt had been applied one year earlier were shorter, had a lower percentage of white clover and a higher percentage of bare ground and balansa clover. In Experiment 2, factorial combinations of salt fertiliser (0, 100 kg NaCl/ha) and seed (no seed, mixture of Caucasian clover (10 kg/ha), *Lotus pendunculatus* (2 kg/ha) and plantain (2 kg/ha)) were applied together to 5 x 10 m plots in September 2004. In April 2005, pastures where salt was applied were shorter, had fewer grass seedheads, a lower percentage of resident grasses and litter but a higher percentage of bare ground and plantain. There were more seedlings of plantain where salt was (6.5 seedlings/m²) than where it was not applied (1.1 seedlings/m²). This small plot work indicates that salt fertiliser application to Na-deficient herbage can enhance grazing intensity. By creating soil disturbance and reducing competition from resident grasses, salt application increased the establishment of oversown seeds of plantain and balansa clover. Establishment of the slow growing species (Caucasian clover, lotus) and the large seeded subterranean clover appeared to be unsuccessful under the conditions of the experiments. Salt application is a tool for pasture management and improvement in the hill/high country that could be used together with other methods such as herbicides, seeding, sub-division, grazing management and fertiliser.

Keywords: grazing, sodium, oversowing, hill country

Introduction

The pasture quality of steep, south-facing slopes in the hill and high country is often poor, with considerable,

tall, rank, laxly-grazed grass, a high proportion of dead material and a low proportion of pasture legume. The establishment of seeds of legumes and herbs oversown onto these sites to improve pasture quality is often negligible due to insufficient grazing to remove the tall competing grass (e.g. cocksfoot, browntop, tall oat grass) vegetation that can inhibit germination and seedling survival (Lambert *et al.* 1985; Awan *et al.* 1993).

Previous survey work indicates that large areas of inland South Island, New Zealand, have pasture sodium (Na) levels in herbage DM less than the 0.07% Na required for a maintenance diet of sheep (Towers & Smith 1983; Aspinall *et al.* 2004). Moreover, Merino sheep are strongly attracted to, and graze intensively, in areas of Na-deficient pasture that have been fertilised with salt (NaCl) (Aspinall *et al.* 2004). These findings raise the possibility of improving pasture quality by using the preference of sheep for applied salt as a tool to intensify grazing and treading on the slopes of Na-deficient inland properties that normally have poor grazing utilisation.

The hypothesis tested in this study was that the destruction of dead material and greater control of grass growth that is achieved with more intense grazing and trampling following salt application would allow existing legumes to flourish and provide conditions that are more conducive to the establishment of oversown seeds.

Materials and methods

Study site

Two experiments were conducted on Na-deficient (<0.03% Na in herbage DM) pasture on south facing slopes (20-35° slope; 540-610 m a.s.l.) at Mt Grand Station, 2 km south east of Hawea Flat, Central Otago. Mt Grand is 150 km from the east coast and 85 km from the west coast, with mean annual rainfall at 600 m a.s.l of 700 mm. The south-facing hill pastures were dominated by three grass species, cocksfoot, tall oat-grass and sweet vernal, with white clover and suckling clover the dominant legumes. Both experiments were located on Arrow stepland soils (Blakemore 1968). Soil samples taken from each site before experiments began and analysed with a MAF quick test showed: Experiment 1, pH = 5.6, Olsen P = 16, Sulphate-S = 10 m.e./100g, Na = 2.0 m.e./100g; and Experiment 2, pH = 6.0, Olsen P = 19, Sulphate-S = 14 m.e./100g, Na = 2.0 m.e./100g. The experiments were carried out in three paddocks where

no salt (neither fertiliser nor blocks) had previously been used.

Oversowing experiments were conducted with five species (balansa clover *Trifolium michelianum*, subterranean clover *T. subterraneum*, Caucasian clover *T. ambiguum*, lotus *Lotus pendunculatus* and plantain *Plantago lanceolata*) that may be agronomically suitable in the Mt Grand environment (Moorhead *et al.* 1994; Stewart & Charlton 2003) but which are potentially difficult to establish.

Experiment 1

This seed oversowing study is a continuation of a fertiliser project established in 2003, with initial results reported by Aspinall *et al.* (2004). The experimental design was three replicates of a 2 x 2 x 2 factorial (eight treatments) laid out in a randomised block design. Two replicates were laid out within Watties (80 ha) and one in Broad Leaf (185 ha) paddocks. The three factors were Na (0 and 150 kg NaCl/ha as coarse 1–4 mm granules), N (0 and 100 kg N/ha as urea) and P+S (0 and 500 kg/ha as 19% sulphur superphosphate). All fertilisers were broadcast by hand on 9 December 2003. Each block consisted of eight 30 x 8 m plots laid out across the slope, with a 30 x 8 m buffer between each plot. To test the residual effects of the fertiliser treatments on oversowing success, half of each plot was oversown with a mixture of balansa (cv. Bolta) and subterranean (Woogenellup) clover at 10 kg/ha per species on 3 March 2004. Lactating Merino ewes were present during the application of fertilisers and within 2 days their intensive grazing and trampling resulted in bare ground approaching 50% versus 20% where no salt was applied. Merino hoggets were present when seed was spread but grazing intensity on all plots was low. Rainfall in autumn after seed spread was 70 mm in March 2004 but only 13 mm in April 2004.

Pasture ground cover of all species was visually estimated in ten 0.1 m² quadrats in each plot on 1–3 December 2004. To provide an indication of grazing intensity, pasture height was measured with a sward stick in 50 locations per plot on 11 November 2004, and 21 January and 14 April 2005. To provide a more precise estimate of balansa clover seedling recruitment following seed application in March 2004, balansa clover inflorescences were counted in five 0.1 m² quadrats in each plot on 28 January 2005.

Experiment 2

The experimental design was six replicates of a 2 x 2 factorial (four treatments) laid out in a randomised block design. All replicates were within Valley (60 ha) paddock. The two factors were Na (0 and 100 kg NaCl/ha as 0.5–1 mm fine granules) and oversown seed (no seed or a

mixture of 10 kg/ha Caucasian clover, cv. Endura, 2 kg/ha *Lotus pendunculatus* cv. Sunrise and 2 kg/ha plantain, cv. Tonic). Both legumes were freshly inoculated with their specific rhizobium strains. Seed and fertilisers were broadcast sown together by hand on 27 September 2004. Each block consisted of four 5 x 10 m plots laid out across the slope, with a 10 x 10 m buffer between each plot. Lactating Merino ewes (280) were present when salt and seed were applied and grazed the salt plots more intensively than elsewhere. Rainfall of 48 mm in September 2004 and 47 mm in October 2004 were less than the average 65 mm per month; but 70 mm in November 2004 and 155 mm in December 2004 exceeded expectations.

Pasture ground cover of all species was visually estimated and seedlings of sown species were counted in ten 0.1 m² quadrats in each plot on 14 April 2005. To provide an indication of grazing intensity, pasture height was measured with a sward stick in 50 locations per plot on 25 November 2004 and 28 January 2005, and seed heads of cocksfoot, tall oat grass and sweet vernal were counted in six 0.1 m² quadrats in each plot on 31 November 2004 and 16 April 2005.

Data from both experiments were analysed by factorial ANOVA. Percentage composition and count data were arcsine square-root transformed and square-root transformed, respectively, before ANOVA.

Results

Experiment 1

When sampled 12 months after the start of the experiment, salt application resulted in pastures that were shorter and with a lower percentage of white clover and resident annual legumes than where no salt was applied. Salt application also resulted in pastures with a higher percentage of bare ground, balansa clover and dicot weeds than where no salt was applied (Table 1). There were more balansa clover inflorescences where salt was applied. Urea application resulted in pastures with a higher percentage of resident grasses and a lower percentage of balansa and resident annual legumes than where no urea was applied (Table 1). Superphosphate application resulted in a higher percentage of white clover (Table 1). There were no significant interactions between salt, nitrogen and superphosphate application.

Experiment 2

Salt application resulted in pastures that were shorter, had fewer grass seed heads and a lower percentage of resident grasses and litter than where no salt was applied (Table 2). The percentage of bare ground and plantain was greater where salt was applied. Seedlings of plantain, but not Caucasian clover or lotus, were found in counts conducted on 14 April 2004. Plantain seedlings were

Table 1 Pasture composition (%) on 1 December 2004, pasture height (cm), and balansa inflorescences/m² after the application of salt (NaCl), nitrogen (urea) and sulphur superphosphate on 9 December 2003. Significance of main effects denoted by * P<0.05, ** P<0.01, ns non significant.

	Salt (kg/ha)			Nitrogen (kg/ha)			Superphosphate (kg/ha)		
	0	150		0	100		0	500	
Pasture composition (%)									
Resident grasses	53.0	53.0	ns	50.1	56.0	*	54.8	51.3	ns
White clover	14.3	10.9	**	12.3	12.8	ns	11.5	13.6	*
Subterranean clover	0.8	1.0	ns	1.1	0.7	ns	1.0	0.8	ns
Balansa clover	1.4	2.3	*	2.3	1.3	*	1.7	2.0	ns
Resident annual legumes	12.1	10.5	*	12.4	10.2	*	11.6	11.0	ns
Litter	9.5	8.0	ns	8.9	8.6	ns	8.8	8.6	ns
Dicot weeds	4.5	5.8	*	5.3	5.0	ns	5.3	5.0	ns
Bare ground	4.4	8.5	*	7.5	5.4	ns	5.2	7.7	ns
Pasture height (cm)									
11 Nov 04	10.7	8.9	*	9.6	10.1	ns	9.8	9.8	ns
21 Jan 05	19.4	16.4	*	17.4	18.6	ns	18.9	16.9	ns
14 Apr 05	20.1	19.0	ns	18.4	20.7	ns	20.9	18.1	ns
Balansa inflorescences/m²									
28 Jan 2005	7.1	25.3	*	22.1	9.6	ns	10.7	21.2	ns

Table 2 Pasture composition (%) on 14 April 2005, pasture height (cm), and seedheads/m² after the application of salt (100 kg NaCl/ha) and seed (plantain, Caucasian clover and lotus) on 27 September 2004. Data are means averaged across plots where seed was and was not sown. Significance of main effects denoted by * P<0.05, ** P<0.01, ns non significant.

	Salt (kg/ha)		
	0	100	
Pasture composition (%)			
Resident grasses	62.5	48.5	**
White clover	9.4	12.2	ns
Annual legumes	1.0	1.5	ns
Dicot weeds	3.3	3.7	ns
Plantain	0.1	2.3	*
Litter	19.1	9.9	**
Bare ground	4.6	22.4	**
Pasture height (cm)			
25 Nov 04	10.9	5.8	*
28 Jan 05	23.6	6.8	*
Seedheads/m²			
31 Nov 04 Cocksfoot	43	10	**
Tall oat	17	3	*
Sweet vernal	315	89	**
16 Apr 05 Cocksfoot	65	1	**
Tall oat	52	1	**
Sweet vernal	117	10	**

only found on sown plots, with the number of plantain seedlings greater where salt was applied (6.5 seedlings/m²) than where it was not applied (1.1 seedlings/m²) (salt x seed interaction, P<0.05).

Discussion

Seedling recruitment of legumes and herbs from natural reseeding or following oversowing in pastures is often

limited, particularly in the hill and high country, due to insufficient grazing to remove competing vegetation (Lambert *et al.* 1985; Edwards *et al.* 2005). This study provided some evidence that this limitation could be overcome by the application of salt to enhance grazing intensity. Both balansa clover in Experiment 1 and plantain in Experiment 2 showed greater seedling establishment where salt was applied. The most likely explanation for

the increased establishment with salt application is the greater percentage of bare ground and reduced competition from resident grass species (cocksfoot and tall oat-grass) associated with more intense grazing. This may have enhanced either germination or seedling survival, although measurements were not taken to distinguish between these two processes in this study. Previous studies have shown that seedling establishment of plantain increases as the proportion of bare ground increases (Edwards *et al.* 2005). The result is also consistent with previous hill and high country work that shows intensive 'hoof and tooth' grazing opens up pastures and enhances establishment (MacFarlane & Bonish 1986; Lambert *et al.* 1985). Thus, it appears that the application of salt increased stocking rate within a specific area of a large block without the requirement for fencing or shepherding stock, and so may be a tool that could be used together with other methods such as herbicides, subdivision and fertilisers to improve oversowing success.

No seedlings of lotus or Caucasian clover were found and there was no detectable increase in subterranean clover cover following oversowing. The poor establishment of lotus and Caucasian clover is consistent with previous studies (Brock & Charlton 1978; Moorhead *et al.* 1994; Black *et al.* 2002), where establishment is slow, even under cultivated conditions where competition is minimised. The poor establishment does not reflect low seed viability as this was >87% in all species in laboratory tests. Alternative explanations include the high thermal time requirements for seedling development of these species (Black *et al.* 2002), desiccation from lack of cover and poor radicle entry due to seeds remaining on the soil surface (Campbell & Swain 1973). Drilling has been shown to significantly increase Caucasian clover seedling establishment above that of oversowing (Moorhead *et al.* 1994), while subterranean clover seeds have higher germination when buried (Yates 1957).

Aspinall *et al.* (2004) reported small areas treated with 100 kg NaCl/ha at the start of Experiment 1 attracted extreme grazing and trampling by Merino ewes, with up to 50% bare ground created within 1 week of salt application. Similar responses were also observed in Experiment 2, where sheep were observed grazing intensively on plots within 48 h of salt application; pastures remained shorter over the next 7 months and had fewer grass seed heads. Measurements reported here from Experiment 1 indicated that the effects of salt application on grazing preference may be long lasting, with bare ground percentage higher and pasture height shorter 12 months after salt application. This sustained preference for areas where salt was applied may reflect two processes. First, it may be due to the preference of sheep for herbage with elevated Na content. Herbage Na

was not measured after 12 months in Experiment 1. However, in a parallel experiment (Gillespie 2006) at the same site in which 150 kg/ha of salt was spread at the same time as in Experiment 1, the Na content of cocksfoot herbage 12 months after salt application was greater where salt was applied (0.08% of DM) than where it was not applied (0.03% DM). Second, it may be due to preference for short vegetative material lacking dead and reproductive material that was created by the initial salt application and intensive grazing.

The percentage of white clover in the pasture was lower where salt was applied in Experiment 1 and unaffected by salt application in Experiment 2. Thus, there was no evidence to support the idea that the shorter pasture, lower percentage of resident grasses and higher percentage of bare ground resulting from salt application allowed white clover stolons to expand into vacant space and increase in abundance. The observed response could be due to treading damage of stolons soon after salt application combined with preferential grazing of short developing clover stolons.

The effect of nitrogen and superphosphate fertiliser on the vegetation was small compared to salt application. The initial nitrogen fertiliser boost to grass growth remained in Experiment 1. The 6% extra ground cover of resident grasses in the nitrogen-fertilised plots may have caused the small reduction in annual clover cover. While there was a small increase in white clover cover with the addition of sulphur superphosphate, there was no response in the annual clovers.

ACKNOWLEDGEMENTS

We thank: Dominion Salt Ltd and the Miss E A, Hellaby Indigenous Grassland Trust for financial support for the project; Gordon Meikle, Mt Grand Farm Manager, and Tony Whatman, Lincoln University Director of Farms for their advice and technical support with stock.

REFERENCES

- Awan, M.H.; Kemp, P.D.; Choudhary, M.A.; Barker, D.J. 1993. Pasture legume establishment from oversowing in drought-prone hill country. *Proceedings of the New Zealand Grassland Association* 55: 101-104.
- Aspinall, R.; Mandaluniz, N.; Hight, L.J.; Lucas, R.J. 2004. Sodium deficiency in Canterbury and Central Otago sheep pastures. *Proceedings of the New Zealand Grassland Association* 66: 227-232.
- Blakemore, L.C. 1968. General survey of the soils of South Island, New Zealand. *In: New Zealand Soil Bureau Bulletin* 27.
- Black, A.D.; Moot, D.J.; Lucas, R.J. 2002. Seedling development and growth of white clover, Caucasian clover and perennial ryegrass grown in field and

- controlled environments. *Proceedings of New Zealand Grassland Association* 64: 197-204.
- Brock, J.L.; Charlton, J.F.L. 1978. *Lotus pedunculatus* establishment in intensive farming. *Proceedings of New Zealand Grassland Association* 39: 121-129.
- Campbell, M.H.; Swain, F.G. 1973. Effect of strength, tith and heterogeneity of soil surface on the radicle-entry of surface-sown seeds. *Journal of the British Grasslands Society* 28: 41-50.
- Edwards, G.R.; Hay, M.J.M.; Brock, J.L. 2005. Seedling recruitment dynamics of forage and weed species under continuous and rotational sheep grazing in a temperate New Zealand pasture. *Grass and Forage Science* 60: 186-199.
- Gillespie, B.J. 2006. Modifying merino grazing and pasture composition in the high country by salt application. MAgSci Thesis. Lincoln University.
- Lambert, M.G.; Rhodes, A.P.; Barker, D.J.; Bircham, J.S. 1985. Establishing and managing improved plants in hill country. *Using herbage cultivars. Grassland Research and Practice Series No. 3*: 31-34.
- Macfarlane, M.J.; Bonish, P.M. 1986. Oversowing white clover into cleared and unimproved North Island hill country. The role of management, fertiliser, inoculation, pelleting and resident rhizobia. *Proceedings of New Zealand Grassland Association* 47: 43-51.
- Moorhead, A.J.E.; White, J.G.H.; Jarvis, P.; Lucas, R.J.; Sedcole, J.R. 1994. Effect of sowing method and fertiliser application on establishment and first season growth of Caucasian clover. *Proceedings of the New Zealand Grassland Association* 56: 91-95.
- Stewart, A.V.; Charlton, J.F.L. 2003. Pasture and forage plants for New Zealand. New Zealand Grassland Association and New Zealand Grassland Trust. Grassland Research and Practice Series No 8. 96 pp.
- Towers, N. R.; and Smith, G. S. 1983. Sodium. The mineral requirements of grazing ruminants. New Zealand Society of Animal Production. pp 115-124 *In: Occasional Publication No. 9.*
- Yates, J.J. 1957. Seed-setting in subterranean clover (*Trifolium subterraneum* L.). 1. The importance of the micro-environment. *Australian Journal of Agricultural Research* 8: 433-443.