

Nitrogen fixation by caucasian clover and white clover in irrigated ryegrass pastures

K.H. WIDDUP¹, R.G. PURVES², A.D. BLACK³, P. JARVIS³ and R.J. LUCAS³

¹AgResearch, PO Box 60, Lincoln

²148 Fisher Avenue, Beckenham, Christchurch

³Soil, Plant and Ecological Sciences Division, PO Box 84, Lincoln University
keith.widdup@agresearch.co.nz

Abstract

The N₂ fixation ability of caucasian clover was compared with that of white clover in irrigated ryegrass pastures over years 2 and 3 of a grazing experiment, using the ¹⁵N enrichment technique. 'Endura' caucasian clover was inoculated with the specific *Rhizobium* strain ICC148. The N concentration in clover herbage and the proportion of clover N derived from N₂ fixation (P_N) were similar for both clovers at averages of 4.6%N and 50–60% respectively over the 2 years. The amount of N₂ fixed per hectare was directly related to the amount of clover dry matter (DM) produced by the two clover species. Caucasian clover produced four times the DM yield of white clover in year 2 (5400 cf. 1450 kg DM/ha) and four times the amount of N₂ fixed in herbage (136 cf. 36 kg N/ha). In year 3, caucasian clover produced 50% more clover DM (3450 cf. 2370 kg DM/ha) and N₂ fixed (98 cf. 66 kg N/ha) than white clover. The increased N input from caucasian clover increased grass %N and N uptake from soil in caucasian clover pastures resulting in higher total pasture production compared with white clover pastures (15.7 cf. 14.2 t DM/ha) by year 3. In this study, caucasian clover demonstrated greater potential than white clover to meet the N demands of high-yielding perennial ryegrass in an intensive pastoral system.

Keywords: clover content, *Lolium perenne*, nitrogen fixation, pasture production, *Trifolium ambiguum*, *T. repens*

Introduction

Biological nitrogen (N₂) fixation by legumes is both ecologically and economically important to pastoral farming in New Zealand (Walker 1995). White clover (*Trifolium repens*) is the predominant legume used and studies have shown that annual N₂ fixation levels averaged 185 kg N/ha/year over 10 New Zealand sites (Hoglund *et al.* 1979). At Kirwee, Canterbury, dryland and irrigated pastures fixed 120 and 190 kg N/ha/year

respectively (Crush 1979). New Zealand's white clover-based pastures, however, are not fixing sufficient nitrogen to support the demands of high-yielding grasses nor to provide enough quality feed for maximum animal production (Caradus *et al.* 1996; Chapman *et al.* 1995; Clark & Harris 1995).

Caucasian clover (*Trifolium ambiguum*) has been demonstrated recently to have potential in lowland, intensive farming systems where it has consistently produced greater total legume yields than white clover in mixed pastures (Black *et al.* 2000; Moss *et al.* 1996; Watson *et al.* 1996). The performance of caucasian clover improved after the selection of the specific *Rhizobium* strain ICC148. Inoculation with this strain resulted in improved establishment, more effective nodules and greater N₂ fixation especially in high country pastures (Pryor *et al.* 1998). The grazing experiment established at Lincoln University in December 1996 (Black *et al.* 2000) to compare caucasian- and white clover-based ryegrass swards provided the resource to measure the N₂ fixation ability of caucasian clover and compare it with that of white clover.

Material and methods

The layout and management of the grazing experiment at Lincoln University was described by Black *et al.* (2000). The experiment was sown in December 1996 with either 'Endura' caucasian clover or 'Demand' white clover at high or low soil fertility with eight replicates. The 'Endura' caucasian clover seed was inoculated with the specific *Rhizobium* strain ICC148 (Pryor *et al.* 1998). 'Ruanui' zero endophyte ryegrass was direct-drilled into the pure clover swards in March 1997. The N₂ fixation study was initiated in spring 1998. It used five replications of the two clover treatments under high soil fertility (Olsen P of 22). The experiment was irrigated over the summer period (December–March) of the 2 years.

The ¹⁵N enrichment method (Ledgard *et al.* 1985) was used to measure N₂ fixation. Commencing in October 1998, labelled ammonium sulphate (40 atom% ¹⁵N) at 0.05 g N/m² was applied to 1.45 m² of pasture in the 10 plots (two clovers x five replicates) and enclosure

cages were placed over the treated areas. The pasture in the treated areas was selected to ensure that adequate populations of the treatment clover species were present. In the caucasian clover plots, ^{15}N was applied to areas with minimal amounts of volunteer white clover. A 0.25 m² quadrat was cut from each treated area approximately every 4 weeks and sub-samples from these cuts were dissected into clover and grass components before drying to determine dry matter (DM) yield and botanical composition. Separate additional sub-samples of the clover and grass herbage were dried and ground for %N and ^{15}N analysis using a mass spectrometer (Anca 20-20 stable isotope analyser). After every second cut, ^{15}N was re-applied to all 10 plots. The ^{15}N treated areas were not moved in 1998/1999 but new areas were selected and treated with ^{15}N every second cut (8-weekly) in 1999/2000. The plots were sampled six times between November 1998 and June 1999 (year 2 of the grazing experiment) and then eight times between October 1999 and June 2000 (year 3).

The proportion (%) of N fixed by the clovers from atmospheric N₂ (P_N) was calculated by:

$$P_N = 100 \times \frac{(\text{atoms}\%^{15}\text{N}_{\text{grass}} - \text{atoms}\%^{15}\text{N}_{\text{clover}})}{(\text{atoms}\%^{15}\text{N}_{\text{grass}} - B)}$$

where B=0.3663, the atoms% ^{15}N of atmospheric N₂.

The amount of fixed N (kg/ha) in the cut clover herbage was then given by:

$$\text{Fixed N} = \text{clover DM (kg/ha)} \times \text{clover}\%N/100 \times P_N/100$$

Results

Mean herbage %N, P_N and N₂ fixation

Mean %N in clover herbage was similar for both caucasian and white clover during 1998/1999 and 1999/2000 (Table 1). The proportion of the N derived from N₂ fixation (P_N) was also similar for caucasian and white clover. In both clover species, N₂ fixation contributed an average of between 50% and 60% of their herbage N over the 2 years (Table 1). The P_N was highest in late spring–summer at about 75% and lowest in late autumn at 45% for both clovers.

In year 2, caucasian clover fixed more N than white clover throughout the whole growing season (Table 1,

Table 1 Mean clover and grass herbage %N, proportion of N fixed by clover (P_N) and amount of fixed N in caucasian and white clover herbage during year 2 (November 1998 to June 1999) and year 3 (October 1999 to June 2000) with perennial ryegrass in high fertility soil under irrigation.

Treatments	Mean %N		P _N (%)	Fixed N in clover Herbage (kg N/ha/yr)	Estimate of total fixed N [#] (kg N/ha/yr)
	Clover	Grass			
----- Year 2 -----					
Caucasian clover	4.61	3.69	55.6	136	232
White clover	4.55	3.38	51.0	36	61
Isd (5%)	0.15ns	0.14*	6.4 ns	13.4*	
----- Year 3 -----					
Caucasian clover	4.59	3.46	56.7	98	167
White clover	4.59	3.12	60.7	66	112
Isd (5%)	0.11ns	0.13*	5.6 ns	7.7*	

[#] Based on 1.7 x fixed N in legume herbage
ns = not significantly different at P>0.05
* = significant at P<0.05

Figure 1). In year 3, the N fixed by caucasian clover was greater in the summer period but similar to white clover in the cooler autumn period (Figure 1). It is likely that the low December values in year 3 were owing to low soil moisture levels, as irrigation was not initiated on the pastures until mid-December. Mean %N in grass herbage was significantly higher in the grass growing with caucasian clover than in the white clover based pastures during both years (Table 1).

Pasture DM production


In 1998/1999, the caucasian- and white clover-based pastures produced similar total DM (grass plus clover), however, clover DM production was greater from caucasian than white clover pastures (Table 2). DM production from the associated grass in caucasian clover pastures was lower than from white clover pastures (Table 2).

Table 2 Total pasture, clover and grass production from caucasian- and white clover-based swards growing with perennial ryegrass, in high fertility soil, under irrigation, during year 2 (November 1998 to June 1999) and year 3 (October 1999 to June 2000).

Treatments	Clover content (%)	DM production (kg DM/ha/yr)		
		Clover	Grass	Total pasture
----- Year 2 -----				
Caucasian clover	52.8	5400	4440	10,080
White clover	15.1	1450	7440	9160
Isd (5%)	21.8*	2540*	2950*	2190ns
----- Year 3 -----				
Caucasian clover	21.9	3450	11,970	15,740
White clover	16.8	2370	11,830	14,200
Isd (5%)	4.8*	940*	439ns	918*

ns = not significantly different at P>0.05
* = significant at P<0.05

Figure 1 N fixed in caucasian and white clover herbage growing with perennial ryegrass in high fertility soil under irrigation during 1998/1999 and 1999/2000 (ns = non-significant at $P > 0.05$ and * = significant at $P < 0.05$).



In 1999/2000, the caucasian clover-based pastures produced greater total DM compared with pastures sown with white clover (Table 2). The grass DM were similar from both pastures, but clover DM was greater from caucasian clover than white clover pastures (22% cf. 17% clover) resulting in the greater total pasture production. Volunteer white clover produced 340 kg DM/ha (10% of the total clover yield) in the caucasian clover pastures in year 3.

Discussion

Caucasian clover and white clover had similar proportions of fixed N_2 (P_N) in their herbage, which indicates that both clovers have a similar ability to fix N_2 in the presence of perennial ryegrass, under irrigation and in high fertility soil. The amount of N_2 fixed per hectare was directly related to the amount of DM produced by the two clover species. Caucasian clover produced four times more DM than white clover in year 2 (5400 cf. 1450 kg DM/ha) and this was associated with four times the amount of N fixed (136 cf. 36 kg N/ha). In year 3, caucasian clover produced 50% more DM than white clover; the amount of N fixed (kg N/ha/yr) from N_2 fixation in the caucasian clover pastures

was therefore 50% greater than that in the white clover pastures.

Mean %N in clover herbage was similar for both clover species and was constant throughout the year, but the clovers showed variation in P_N across seasons of the year. The P_N was highest in late spring–summer (75%) and lowest in autumn (45%) for both clover species. Ledgard *et al.* (2001) attributed lower P_N levels over autumn and early spring to low temperatures reducing clover growth, and/or elevated soil inorganic N during these periods. The level of soil inorganic N has a major modifying action on clover N_2 fixation – as soil inorganic N availability increases, N_2 fixation decreases (Hoglund *et al.* 1979). High P_N occurred in late spring–summer when high temperatures increased clover growth rates and soil inorganic N levels are normally lower owing to rapid growth and high N uptake by grass.

The association between caucasian clover and the specific *Rhizobium* strain ICC148 was obviously very effective and productive in this experiment. The only other reported N_2 fixation values from caucasian clover under grazing in New Zealand indicated lower values than white clover (Watson *et al.* 1996), but they suggested the acetylene reduction technique and shallow

(75 mm) depth of sampling used were not appropriate for caucasian clover. In addition, the *Rhizobium* strain CC283B used in that experiment has since been surpassed by the new strain ICC148 with improved N₂ fixation ability (Pryor *et al.* 1998).

The ¹⁵N enrichment method provides an accurate estimate of the amount of fixed N in the harvested clover herbage. However, this is likely to be an underestimate of total N fixed because it does not account for fixed N in clover stolons, rhizomes and roots. Jørgensen & Ledgard (1997) showed the amount of fixed N in stolons and roots of white clover averaged 70% of that in clover herbage. Therefore, a 1.7 multiplying factor was used to give an estimate of the total N fixed. Using this conversion factor, the white clover-based pastures produced 61 and 112 kg N/ha in years 2 and 3 respectively while the caucasian clover pastures produced 232 and 167 kg N/ha (Table 1).

This method of estimating the total N fixed is likely to underestimate the value even more for caucasian clover as its shoot:root ratio is 30:70 compared with the 60:40 ratio for white clover (Widdup *et al.* 1998). If the same principles were applied to caucasian clover, the multiplying factor would be 3.3. However, this is likely to be an overestimate as much of the rhizome tissue is a storage organ and only the N in the newly grown rhizome tissue would reflect the N₂ fixation during any regrowth period. Further studies are required to provide estimates of the fixed N in the above-ground herbage compared with below-ground rhizome/root tissues of caucasian clover.

Many New Zealand white clover-based pastures have average clover contents less than 20% (Ettema & Ledgard 1992; Moss 1987). This is generally considered insufficient to support the N requirements of high-yielding grasses nor to provide enough quality feed for maximum animal performance (Caradus *et al.* 1996; Chapman *et al.* 1995; Clark & Harris 1995). Thomas (1992) estimated clover contents of 20–45% could provide the N requirements of a sustainable and productive pasture. Caucasian clover has shown potential to approach these target clover contents when in association with aggressive grass species in irrigated (Black *et al.* 2000; Moss *et al.* 1996) and dryland (Black & Lucas 2000; Watson *et al.* 1996) environments. In this experiment, caucasian clover pastures had a higher clover content and fixed more N₂ than white clover pastures during both years. Caucasian clover pastures did, however, have exceptionally high clover contents in year 2, which can be partly explained by the management of the N₂ fixation plots. Four-weekly cutting with no excreta return for 8 months accentuated the clover component and depressed the grass component. Even though both clover species

received the same cutting management, caucasian clover displayed greater growth potential than white clover under the repeated cutting regime.

In year 3, ¹⁵N was applied to a new caged area every 8 weeks, a management allowing greater exposure of plots to normal grazing and excreta return; the clover contents were lower than in year 2. The lower clover contents were also attributed to a heavy infestation of grass grub (*Costelytra zealandica*) in autumn 1999 (averaged 180 grubs/m²) which greatly reduced the shoot and root material of both clover species (Trevor Jackson, pers. comm.). In spring 1999, white clover had recovered from surviving stolon pieces and new seedlings whereas the recovery of caucasian clover depended solely on new rhizome development from surviving rhizomes and tap-roots. Despite this disadvantage, caucasian clover produced significantly more clover herbage than white clover in the third year.

High N fixation by legumes in mixed pastures leads to greater N input into soil organic N resulting in increased N availability to associated grasses. In this study, the grass %N and the total soil N uptake was greatest in caucasian clover-based pastures by year 3 (Table 1 and 3) which resulted in higher total pasture production during year 3 (Table 2). This response needs to be monitored further at the Lincoln University grazing experiment to determine whether caucasian clover can maintain higher clover and total production than white clover in an environment with high soil N.

Table 3 Total Soil N uptake (kg N/ha/year) in caucasian clover- and white clover-based pastures during year 2 (November 1998 to June 1999) and year 3 (October 1999 to June 2000).

Treatments	Soil N uptake (kg N/ha/yr)		
	Grass N	Legume N [#]	Total Soil N
	----- Year 2 -----		
Caucasian clover	167	111	276
White clover	252	32	281
lsd (5%)	117ns	52*	87ns
	----- Year 3 -----		
Caucasian clover	414	69	483
White clover	369	43	412
lsd (5%)	17*	18*	29*

[#] Legume N uptake from soil = Legume DM * %N * (1-P_w/100)

ns = not significantly different at P>0.05

* = significant at P<0.05

Conclusions

1. Both caucasian clover and white clover had similar proportions of fixed N₂ in their herbage. This indicates that both clovers had a similar N-fixing ability in the presence of perennial ryegrass, irrigation and high fertility soil.

2. The amount of N fixed per hectare by caucasian and white clover was directly related to the herbage DM produced by the two clover species. Caucasian clover produced four times the DM and total N fixed of white clover in year 2 of the grazing experiment and 50% more DM and N fixed in year 3. The total N fixed ranged from 167–232 kg N/ha for caucasian clover and 61–112 kg N/ha for white clover.
3. In the caucasian clover-based pastures, the increased N input from the clover had led to greater soil N uptake by the associated grasses by year 3 resulting in higher total pasture production. Further study is required to better quantify soil N changes and changes in total pasture production over time.

ACKNOWLEDGEMENTS

The authors acknowledge funding from the Struthers Trust for the development of the grazing experiment at Lincoln University and FRST funding for provision of ^{15}N and N analyses.

The assistance of Bruce Pownall for the supply of ewe lambs and maintaining animal health, Keith Pollock and Jo Amyes for field assistance and Stuart Larsen and Roger Cresswell for conducting the N analyses are thanked. We thank the C. Alma Baker and Struthers Trusts for providing A.D. Black with financial support from post-graduate scholarships.

REFERENCES

- Black, A.D.; Pollock, K.M.; Lucas, R.J.; Amyes, J.M.; Pownall, D.B.; Sedcole, J.R. 2000. Caucasian clover/ryegrass produced more legume than white clover/ryegrass pastures in a grazed comparison. *Proceedings of the New Zealand Grassland Association* 62: 69–74.
- Black, A.D.; Lucas, R.J. 2000. Caucasian clover was more productive than white clover in grass mixtures under drought conditions. *Proceedings of the New Zealand Grassland Association* 62: 183–188.
- Caradus, J.R.; Harris, S.L.; Johnson, R.J. 1996. Increased clover content for increased milk production. *Proceedings of the Ruakura Farmers' Conference* 48: 42–49.
- Chapman, D.F.; Parsons, A.J.; Schwinning, S. 1995. Management of clover in grazed pastures: expectations, limitations and opportunities. *White clover: New Zealand's Competitive Edge. Agronomy Society of New Zealand Special Publication 11/Grassland Research and Practice Series* 6: 55–64.
- Clark, D.A.; Harris, S.L. 1995. White clover or nitrogen fertiliser for dairying? *White clover: New Zealand's Competitive Edge. Agronomy Society of New Zealand Special Publication 11/Grassland Research and Practice Series* 6: 107–114.
- Crush, J.R. 1979. Nitrogen fixation in pasture. IX. Canterbury Plains, Kirwee. *New Zealand Journal of Experimental Agriculture* 7: 35–38.
- Ettema, P.J.; Ledgard, S.F. 1992. Getting the best out of white clover. *Proceedings of the Ruakura Farmers' Conference* 44: 72–76.
- Hoglund, J.H.; Crush, J.R.; Brock, J.L.; Ball, R. 1979. Nitrogen fixation in pasture. XII. General Discussion. *New Zealand Journal of Experimental Agriculture* 7: 45–51.
- Jørgensen, F.V.; Ledgard, S.F. 1997. Contribution from stolons and roots to estimates of the total amount of N_2 fixed by white clover. *Annals of Botany* 80: 641–648.
- Ledgard, S.F.; Simpson, J.R.; Freney, J.R.; Bergersen, F.J. 1985. Field evaluation of ^{15}N techniques for estimating nitrogen fixation in legume-grass associations. *Australian Journal of Agricultural Research* 36: 663–676.
- Ledgard, S.F.; Sprosen, M.S.; Penno, J.W.; Rajendram, G.S. 2001. Nitrogen fixation by white clover in pastures grazed by dairy cows: Temporal variation and effects of nitrogen fertilization. *Plant and Soil* 229: 177–187.
- Moss, R.A. 1987. Pasture and animal performances on an irrigated sheep production farmlet. *Winchmore Irrigation Research Station Technical Report* 24: 4–7.
- Moss, R.A.; Burton, R.N.; Allan, B.E. 1996. Productivity of Caucasian clover based pastures under irrigation. *Proceedings of the New Zealand Grassland Association* 58: 177–181.
- Pryor, H.N.; Lowther, W.L.; McIntyre, H.J.; Ronson, C.W. 1998. An inoculant Rhizobium strain for improved establishment and growth of hexaploid Caucasian clover. *New Zealand Journal of Agricultural Research* 41: 179–189.
- Thomas, R.J. 1992. The role of legumes in the nitrogen cycle of productive and sustainable pastures. *Grass and Forage Science* 47: 133–142.
- Walker, T.W. 1995. The value of N-fixation to pastoral agriculture in New Zealand. *White clover: New Zealand's Competitive Edge. Agronomy Society of New Zealand Special Publication 11/Grassland Research and Practice Series* 6: 115–118.
- Watson, R.N.; Neville, F.J.; Bell, N.L.; Harris, S.L. 1996. Caucasian clover as a pasture legume for dryland dairying in the coastal Bay of Plenty. *Proceedings of the New Zealand Grassland Association* 58: 183–188.
- Widdup, K.H.; Knight, T.L.; Waters, C.J. 1998. Genetic variation for rate of establishment in caucasian clover. *Proceedings of the New Zealand Grassland Association* 60: 213–217.

