

**CENTRE FOR
COMPUTING AND BIOMETRICS**

**Effect of intraruminal infusion of
potassium on the site of magnesium
absorption within the digestive tract
in sheep**


D.E. Dalley, P. Isherwood, A.R. Sykes and A.B. Robson

Research Report No:96/10
October 1996

RESEARCH
REPORT

ISSN 1173-8405

LINCOLN
UNIVERSITY
Te Whare Wānaka O Aoraki



Centre for Computing and Biometrics

The Centre for Computing and Biometrics (CCB) has both an academic (teaching and research) role and a computer services role. The academic section teaches subjects leading to a Bachelor of Applied Computing degree and a computing major in the BCM degree. In addition it contributes computing, statistics and mathematics subjects to a wide range of other Lincoln University degrees. The CCB is also strongly involved in postgraduate teaching leading to honours, masters and PhD degrees. The department has active research interests in modelling and simulation, applied statistics and statistical consulting, end user computing, computer assisted learning, networking, geometric modelling, visualisation, databases and information sharing.

The Computer Services section provides and supports the computer facilities used throughout Lincoln University for teaching, research and administration. It is also responsible for the telecommunications services of the University.

Research Report Editors

Every paper appearing in this series has undergone editorial review within the Centre for Computing and Biometrics. Current members of the editorial panel are

Dr Alan McKinnon	Dr Keith Unsworth
Dr Bill Rosenberg	Dr Don Kulasiri
Dr Clare Churcher	Mr Kenneth Choo
Dr Jim Young	

The views expressed in this paper are not necessarily the same as those held by members of the editorial panel. The accuracy of the information presented in this paper is the sole responsibility of the authors.

Copyright

Copyright remains with the authors. Unless otherwise stated, permission to copy for research or teaching purposes is granted on the condition that the authors and the series are given due acknowledgement. Reproduction in any form for purposes other than research or teaching is forbidden unless prior written permission has been obtained from the authors.

Correspondence

This paper represents work to date and may not necessarily form the basis for the authors' final conclusions relating to this topic. It is likely, however, that the paper will appear in some form in a journal or in conference proceedings in the near future. The authors would be pleased to receive correspondence in connection with any of the issues raised in this paper. Please contact the authors either by email or by writing to the address below.

Any correspondence concerning the series should be sent to:

The Editor
Centre for Computing and Biometrics
PO Box 84
Lincoln University
Canterbury, NEW ZEALAND
Email: computing@lincoln.ac.nz

Effect of intraruminal infusion of potassium on the site of magnesium absorption within the digestive tract in sheep

D.E. Dalley *, P. Isherwood, A.R. Sykes, and A.B. Robson#

Animal and Veterinary Sciences Group, Lincoln University,
Canterbury, New Zealand

*Present address : Agriculture Victoria, Ellinbank, RMD 2460,
Warragul, Victoria, 3820, Australia

#Centre for Computing and Biometrics, Lincoln University,
Canterbury, New Zealand

SUMMARY

Sixteen 2-year old female sheep were fitted with rumen and duodenal cannulae at Johnstone Memorial Laboratory, Lincoln University during 1989-1990. They were offered, at 2 hourly intervals, a pelleted concentrate diet (900 g/day) and chaffed lucerne hay (100 g/day). In a split plot experiment they were infused, intraruminally and at four rates, with potassium (providing 16, 26, 36 or 46 g K/kg food DM/day) and magnesium (providing 1.3, 1.8, 2.3 or 3.1 g Mg/kg food DM/day) within a latin square design and with the liquid and solid phase markers ⁵¹Chromium EDTA and ¹⁴¹Cerium chloride. Net absorption of Mg before and after the duodenum was estimated from dietary intake, duodenal flow and urinary and faecal excretion of Mg.

Increasing K intake resulted in a decline in net absorption of Mg from the entire digestive tract, supporting data in the literature. Increasing K intake from 16 to 46 g/kg DM decreased urinary Mg excretion by between 0.14 and 0.30 g/day, the extent of which was independent of the level of Mg intake. At high K intake Mg absorption from the rumen was reduced, the amount absorbed ranging from 0.08 g Mg/day at intakes of 1.3 g Mg/day and 16 g K/kg DM/day to 0.46 g Mg/day at intakes of 3.1 g Mg/day and 16 g K/kg DM/day. However, at high K intake, and when Mg absorption from the rumen was reduced, net Mg absorption from sites distal to the rumen was increased to an extent which suggested compensatory absorption. Increase in K intake was associated with a consistent reduction in plasma Mg concentration which was independent of

Mg intake. Increases in Mg intake resulted in increases in Mg absorption and plasma Mg concentration at all rates of K intake in direct proportion to rate of intake. The reduction in Mg absorption from the rumen at high K intake was associated with an increase (0.3 units) in pH of rumen digesta.

Short title : Effects of K on Mg absorption in sheep

Corresponding author: Dr A.B. Robson, Centre for Computing & Biometrics, PO Box 84, Lincoln University, Canterbury, New Zealand.

INTRODUCTION

It is well established that dietary potassium (K) decreases magnesium (Mg) absorption in ruminants (Newton *et al.* 1972; Tomas & Potter 1976a ; Field & Suttle 1979 ; Greene *et al.* 1983a,b,c ; Martens & Blume 1986). Quantitative data showing the response surfaces of Mg absorption to changes in dietary Mg and K encountered under grazing conditions, suitable for incorporation into a simulation model of Mg metabolism (Robson 1991) are not, however, available.

Little is known about the effects of high K intake on the partitioning of net Mg absorption between sites in the gastrointestinal tract. The rumen is considered to be the major site of Mg absorption (Tomas & Potter 1976a,b ; Field & Munro 1977 ; Gabel & Martens 1985) but there is evidence that absorption can also occur from the large intestine (Grace & MacRae 1972 ; Ben-Ghedalia *et al.* 1975 ; Tomas & Potter 1976b ; Dalley & Sykes 1989). The role of the large intestine in Mg homeostasis, particularly under conditions of depressed ruminal Mg absorption, is not clear.

This trial was designed first to study the interactions between dietary Mg and K within the range of concentrations encountered in forages and second to measure the partitioning of absorption between the rumen and post ruminal sites. A further objective was to establish whether increasing Mg intake would overcome the depression in Mg absorption that occurs at high K intake.

MATERIALS AND METHODS

Sixteen 2-year old female Coopworth sheep, average weight 43 ± 3 kg, were fitted with a rigid permanent cannula (3 cm internal diameter) in the rumen and a T-shaped cannula in the proximal duodenum (c. 50-100 mm distal to the pylorus), at least 2 months prior to experimentation. This surgery was carried out at the Johnstone Memorial Laboratory, Lincoln University during 1989-1990. Normal cannulation techniques were used (Hecker 1974).

Animals were housed indoors in metabolism crates designed for separation and collection of urine and faeces. They were offered 900 g/day of a pelleted diet based on barley grain, barley malt culms and barley straw with a trace

element premix and 100 g/day of chaffed lucerne (*Medicago sativa*) hay, both of which were delivered at 2 h intervals, and had free access to fresh tap water.

Animals, ranked hierarchically by weight, were allocated to one of 4 K treatment groups, 0, 8.9, 17.8 and 26.7 g K/day for K treatments 1 - 4, respectively. Within each K treatment four rates of Mg infusion, 0, 0.5, 1 and 2 g/day, were randomly allocated within a latin square design.

The K and Mg supplements, provided as aqueous solutions of chloride salts, were infused into the rumen through polyvinyl chloride tubing at a rate of 460 ml/day during 12 d periods. Sodium bicarbonate was infused into the rumen, independently of K and Mg, to deliver 0.8 g Na/animal/day in 144 ml water. This was done to protect animals from Na deficiency at high K infusion rates, a situation which occurred in a preliminary trial.

Digesta flow past the duodenal cannula was calculated according to the double marker technique of Faichney (1975) using the ⁵¹chromium (⁵¹Cr) complex of ethylenediaminetetraacetic acid (⁵¹Cr EDTA ; Downs & MacDonald 1964) and ¹⁴¹Ce (cerium chloride in 0.1 M hydrochloric acid (HCl)) as liquid and solid phase markers, respectively. Approximately 40 µCi ⁵¹Cr EDTA and 8 µCi ¹⁴¹Ce were included in the sodium bicarbonate solution during the final 8 days of each infusion period. Samples (50 ml) of duodenal digesta were collected at 4 h intervals and rumen fluid (200 ml) at 8 h intervals during the last 4 days of the infusion period and stored at 4 ° C.

Feed offered and refusals were recorded and subsampled daily. Faeces and urine were collected and measured daily throughout the 12 day infusion period. Urine was acidified to pH 2-3 with 1 M HCl and the faeces homogenized before each were subsampled and stored at -20°C. At 08:30 h throughout the infusion period, 5 ml jugular blood was withdrawn through an indwelling jugular catheter, transferred to blood tubes containing lithium heparin (72 USP units), centrifuged at 1200 g for 15 min and the plasma stored at -20 °C.

Laboratory analysis

Samples of rumen and duodenal digesta were centrifuged at 30 000 g for 30 min and the supernatant removed. Further aliquots (10 ml) of rumen and duodenal digesta were suspended in agar (0.5 g) in counting tubes by heating, taking care to maintain a uniform suspension, until the agar set. Radioactivity in both the whole digesta and supernatant fractions was counted in a universal gamma counter (1282 Compugamma, LKB, Wallac).

Feed, faeces and digesta samples were dried to constant weight at 100 °C. Feed and faeces samples were wet ashed by the technique of Thompson & Blanchflower (1971). Rumen and duodenal digesta samples were ashed at 550 °C for 12h. All digests were taken up for subsequent analysis in 5 % (v:v) HCl. Magnesium in these samples and in urine, plasma and rumen and duodenal supernatant fractions was determined by atomic absorption

spectrophotometry following dilution in 0.1 M HCl containing 2000 ppm strontium. Sodium and potassium in feed, plasma and urine were determined by flame emission spectrophotometry after dilution in a 1000 ppm lithium carbonate solution.

Magnesium solubility was calculated as the proportion of total digesta Mg present in the 30 000 g supernatant fraction.

Statistical analysis

For ruminal and duodenal flow measurements, the balance period was the last 4 days of the infusion period, whereas for urinary and faecal excretion measurements the balance period was the last 8 days of the infusion period. Data collected during the balance period were analysed as a split-plot experiment for main effects and interactions using the anova procedure of GENSTAT 5, release 2.2 (Lawes Agricultural Trust, 1991) to calculate estimated treatment means.

RESULTS

The concentrate diet and lucerne hay contained 890 and 920 g DM/kg fresh weight, respectively and the ration fed had an apparent DM digestibility of 0.72. Individual sheep mean, within-period DM intake ranged from 830 to 893 g/day and was unaffected by treatment. Mineral concentrations in the two feeds are presented in Table 1. Mean dietary Mg, Na and K intakes were 1.3, 0.76 and 15.0 g/day, respectively.

Potassium and magnesium infusion rates

The four K treatments provided total daily K supplies of 15, 24, 33 and 42 g/day, or the equivalent of 16, 26, 36 and 46 g K/kg DM for K treatments 1-4, respectively.

Magnesium infusions supplied between 0 and 1.86 g Mg/day which resulted in total daily Mg supplies of 1.3, 1.8, 2.3 and 3.1 g/day, or the equivalent of 1.5, 2.0, 2.5 and 3.5 g Mg/kg DM for Mg treatments 1-4, respectively.

Urinary Mg excretion

A significant positive relationship ($P < 0.001$) was observed between the rate of Mg intake and urinary Mg excretion (Table 2). Increasing the K supply from 16 to 46 g/kg DM reduced urinary Mg excretion ($P < 0.001$) linearly across all Mg treatments (Table 2). In absolute terms the reduction in urinary Mg excretion was similar for all K infusion rates. However, as a proportion of total urinary excretion the reduction was greatest at low Mg supplies.

Urinary potassium excretion

A significant linear increase in mean urinary K excretion ($P < 0.001$) was observed as K supply increased. Mean urinary K excretion values ranged from 5.8 for K1 to 25.4 g/day for K4 (sem 0.68). Overall, urinary K excretion values, irrespective of Mg supply, represented 38, 55, 58 and 60 % of total daily K supply for K treatments 1-4, respectively. Magnesium supply had no significant effect ($P > 0.05$) on urinary K excretion.

Faecal magnesium excretion

Mean faecal Mg excretion was positively linearly related to Mg ($P < 0.001$) but not K ($P > 0.05$) supply (Table 2). However, a significant carry-over effect was observed, which was still present in further analysis when the data were pooled over the last three days of the balance period.

Plasma magnesium

Significant positive linear relationships ($P = 0.001$) were observed between Mg supply and plasma Mg concentration and linear negative relationships ($P < 0.001$) with K supply (Table 2). In general the effect of K in depressing plasma Mg concentration was least at the lowest rate of supply of Mg, though the effects of high K and low Mg supply on plasma Mg were independent but additive in effect.

Plasma potassium

A significant ($P < 0.01$) linear increase in plasma K concentration was observed as K supply increased from 16 to 46 g/kg DM on all Mg treatments. Mean plasma K concentration was 4.35 (± 0.058) and 4.73 (± 0.063) mmol/l for K1 and K4, respectively. Magnesium supply had no significant effect on plasma K concentration.

Rumen digesta

There was no effect of Mg supply on rumen pH or Mg solubility in rumen digesta and the data have been combined within K treatments and are given in Table 3. There was a significant cubic effect of K supply on digesta pH ($P < 0.05$), with the two highest K treatments having the highest pH. The concentration of Mg in whole rumen contents increased linearly with increase in Mg supply ($P < 0.001$) but not with an increase in K supply ($P > 0.05$). Magnesium concentration in the supernatant fraction of rumen digesta increased with increase in Mg supply ($P < 0.001$), and tended to decrease with increase in K supply.

Duodenal digesta flow

Duodenal digesta flow increased ($P < 0.005$) with increasing K supply from 10.2 during K treatment 1 to 12.3 l/day during K treatment 4 (sem 0.46) and was associated with a non significant increase in rumen volume from 3.1 to 3.7 l for K treatments 1 and 4, respectively. There was a tendency for digesta flow rate to increase with Mg supply, but this increase was not significant.

The flow of Mg in whole duodenal digesta (Table 2) increased linearly with both K ($P < 0.05$) and Mg ($P < 0.001$) supply though the effect of Mg was greater than that of K.

DISCUSSION

The amount of Mg disappearing before the duodenum at the low K intake increased with increasing Mg supply, as would be expected from the work of Martens *et al.* (1978) and McLean *et al.* (1984) which showed a direct relationship between rumen Mg concentration and rate of Mg absorption. In addition, at this level of K supply the rumen appeared to be the major site of Mg absorption, as would be anticipated from the studies of Grace & MacRae (1972), Tomas & Potter (1976b) and Field & Munro (1977). The increase in duodenal Mg flow with increasing K supply (Table 2) further confirms the findings of Tomas & Potter (1976c) which have demonstrated impairment of Mg absorption from the rumen with increasing rumen fluid K concentration.

A significant finding, however, was that despite negligible absorption of Mg from the rumen at high K supply, between 0.2 and 0.7 g/day of Mg was excreted in urine. This must suggest that compensatory absorption was occurring from some site distal to the rumen (Fig. 1) particularly since this could not be furnished by the depression in plasma concentration and skeletal Mg cannot be rapidly mobilized in mature sheep (Terashima *et al.* 1987). The data in Fig. 1 clearly show that not only was less Mg absorbed from the entire digestive tract as K supply increased but the site of net absorption also changed, the amount absorbed from the hindgut increasing significantly. Tomas & Potter (1976b) showed the large intestine to be capable of absorption of substantial quantities of Mg when Mg was supplied by infusion into the ileum rather than the rumen and McLean *et al.* (1984) showed that with increasing rate of Mg infusion increasing proportions of the Mg absorbed were absorbed distal to the pylorus. Furthermore, Grace & MacRae (1972) observed that while only 2 % of net Mg absorption occurred distal to the pylorus in continuously fed sheep, 50 % was absorbed at this site in sheep fed once daily which they interpreted as resulting from a pulse of Mg which overloaded the rumen capacity for transport and became available for absorption distally. The present work suggests that distal sites become more important in K induced reduction in absorption from the rumen (Fig. 1) and could well be the sites from which orally administered Mg supplements are absorbed. Interestingly, Robson *et al.* (1996) found it necessary to assume hindgut absorption of Mg in

order to satisfactorily explain experimental observations from balance trials in development of a computer simulation model of Mg absorption.

The distal site of Mg absorption in the present experiment is most likely to be the large intestine as net secretion of Mg from the small intestine has been reported in many experiments (Ben-Ghedalia *et al.* 1975 ; Tomas & Potter 1976a ; Bown *et al.* 1989) and there are several reports of net Mg absorption from the large intestine (Tomas & Potter 1976a ; McLean *et al.* 1984 ; Dalley & Sykes 1989). Little is known, however, about the mechanism of absorption from the hindgut. The increase in Mg absorption from the large intestine observed by Tomas & Potter (1976b) was associated with a 20 - 25 % increase in Mg entering the large intestine, and in the present work a greater Mg flow and concentration was probably responsible for the observed increase in Mg absorption. Although McLean *et al.* (1984) demonstrated a much lower potential difference between the duodenum and plasma than between rumen and plasma (13.30 and 36.67 mV, respectively), which would facilitate diffusional transport, nothing is known about conditions and transport mechanisms in the hindgut. Clearly it is either not subject to the same depression by digesta K concentration or K concentration in that part of the tract is not significantly affected by K intake. The former might be the more likely since we had indirect evidence, in urinary K excretion, that K content in hindgut digesta would have increased with K supplementation.

The effect of K and Mg on retention of Mg in the body was estimated by multiplying plasma Mg concentration by the estimated volume of the extracellular fluid (ECF) pool (0.15 X weight ; Storry 1961), assuming ECF Mg was in equilibrium with plasma Mg. Despite significant increases in plasma Mg concentration with increasing Mg supply and decreases with increasing K supply, actual quantities of Mg amounted to only 0.1 - 0.3 g, indicating that retention of Mg was small relative to daily throughput of Mg.

Coefficients of absorption for Mg ranged from 0.13 at low Mg and high K intakes to 0.35 at high Mg and low K intakes and span the range anticipated in mature ruminants (ARC 1980). The very low coefficient of absorption observed at high K supplies when combined with a low Mg supply was less than the conservative value of 0.17 used in most calculations of Mg requirement (ARC 1980). The K concentrations used in the present work (16-46 gK/kg DM) represent the range experienced by pastoral ruminants. It may be that the magnitude of the depression in net Mg absorption at high K and low Mg intake is greater than previously thought.

The decrease in Mg absorption from the rumen with increasing K supply was associated with a small increase in rumen digesta pH which depressed Mg concentrations in the liquid phase of digesta. Smith & Horn (1976) demonstrated *in vitro*, using rumen fluid, a decline in Mg solubility with increasing pH, the critical pH for change in solubility being 6.5 to 7.0. Although rumen digesta pH at the low K supply in the present experiment was c. 5.8, at high K supplies it increased to 6.0 and was associated with a decline in Mg

solubility from c. 0.37 to 0.33. The greatest effect of K supply on Mg solubility was observed at low Mg supply where it was 0.42 and 0.29 on treatments with K1 and K4, respectively. Similar effects of K were observed by Greene *et al.* (1983c) and Wylie *et al.* (1985), though these workers used potassium bicarbonate as the K source and change in pH may have been due to the increase in bicarbonate ion concentration.

REFERENCES

Agricultural Research Council (1980). The Nutrient Requirements of Ruminant Livestock. Slough: Commonwealth Agricultural Bureaux, London.

Ben-Ghedalia, D., Tagari, H., Zamwel, S. & Bondi, A. (1975). Solubility and net exchange of calcium, magnesium and phosphorus in digesta flowing along the gut of the sheep. *British Journal of Nutrition* **33**, 87-94.

Bown, M.D., Poppi, D.P. & Sykes, A.R. (1989). The effects of concurrent infection of *Trichostrongylus colubriformis* and *Ostertagia circumcincta* on calcium, phosphorus and magnesium transactions along the digestive tract of lambs. *Journal of Comparative Pathology* **101**, 11-20.

Dalley, D.E. & Sykes, A.R. (1989). Magnesium absorption from the large intestine of sheep. *Proceedings of the New Zealand Society of Animal Production* **49**, 229-232.

Downs, A.M. & MacDonald, I.W. (1964). The chromium-51 complex of ethylenediaminetetraacetic acid as a soluble rumen marker. *British Journal of Nutrition* **18**, 153-162.

Faichney, G.J. (1975). The use of markers to partition digestion within the gastro-intestinal tract of ruminants. In *Digestion and Metabolism in the Ruminant* (Eds I.W. McDonald & A.C.I. Warner). pp277-291 Armidale, NSW, Australia: University of New England Publishing Unit.

Field, A.C. & Munro, C.S. (1977). The effect of site and quantity on the extent of absorption of magnesium into the gastro-intestinal tract of sheep. *Journal of Agricultural Science, Cambridge* **89**, 365-371.

Field, A.C. & Suttle, N.F. (1979). Effect of high potassium and low magnesium intakes on the mineral metabolism of monozygotic twin cows. *Journal of Comparative Pathology* **89**, 431-439.

Gabel, G. & Martens, H. (1985). Magnesium absorption from the rumen of heifers. *Zentralblatt für Veterinärmedizin Reihe A* **32**, 636-639.

Grace, N.D. & MacRae, J.C. (1972). Influence of feeding regimen and protein supplementation on the sites of net absorption of magnesium in sheep. *British Journal of Nutrition* **27**, 51-55.

Greene, L.W., Fontenot, J.P. & Webb, K.E. Jr. (1983a). Effect of dietary potassium on absorption of magnesium and other macro-elements in sheep fed different levels of magnesium. *Journal of Animal Science* **56**, 1208-1212.

Greene, L.W., Webb, K.E.Jr. & Fontenot, J.P. (1983b). Effect of potassium level on site of absorption of magnesium and other macro-elements in sheep. *Journal of Animal Science* **56**, 1214-1221.

Greene, L.W., Fontenot, J.P. & Webb, K.E. Jr. (1983c). Site of magnesium and other macro-mineral absorption in steers fed high levels of potassium. *Journal of Animal Science* **57**, 503-510.

Hecker, J.F. (1974). Experimental Surgery on Small Ruminants. London: Butterworths.

Lawes Agricultural Trust (1991). Genstat 5, Release 2.2, Reference Manual. Oxford: Clarendon Press.

Martens, H. & Blume, I. (1986). Effect of intraruminal sodium and potassium concentrations and of the transmural potential difference on magnesium absorption from the temporarily isolated rumen of sheep. *Quarterly Journal of Experimental Physiology* **72**, 181-188.

Martens, H., Harmeyer, J. & Michael, H. (1978). Magnesium transport by the isolated rumen epithelium in sheep. *Research in Veterinary Science* **24**, 161-168.

McLean, A.F., Buchan, W. & Scott, D. (1984). Magnesium absorption in mature ewes infused intraruminally with magnesium chloride. *British Journal of Nutrition* **52**, 523-527.

Newton, G.L., Fontenot, J.P., Tucker, R.E. & Polan, G.E. (1972). Effects of high dietary potassium intake on the metabolism of sheep. *Journal of Animal Science* **35**, 440-445.

Robson, A.B. (1991). *Modelling of magnesium metabolism in ruminants*. PhD thesis, Lincoln University, New Zealand.

Robson, A.B., Field, A.C., Sykes, A.R. & McKinnon, A.E. (1996) A model of magnesium metabolism in young sheep. Mg absorption and excretion. *Submitted to British Journal of Nutrition*.

Smith, R.H. & Horn, J.P. (1976). Absorption of magnesium labelled with Magnesium-28 from the stomach of the young steer. In *International Symposium on Nuclear Techniques in Animal Production and health, Vienna*. pp. 253-260. Vienna : International Atomic Energy Agency.

Storry, J.E. (1961). Calcium and magnesium contents of various secretions entering the digestive tract of sheep. *Nature* **4782**, 1197-1198.

Terashima, Y., Matsunobu, S., Yanagisawa, Y. & Itoh. H. (1987) Calcium mobilization in hypomagnasemic wethers fed on a low magnesium and/or high potassium diet. *Japanese Journal of Zootechnical Sciences* **59**, 75-81.

Thompson, T.H. & Blanchflower, W.J. (1971). Wet ashing apparatus to prepare biological materials for atomic absorption spectrophotometry. *Laboratory Practice* **20**, 859-861.

Tomas, F.M. & Potter, B.J. (1976a). The site of magnesium absorption from the ruminant stomach. *British Journal of Nutrition* **36**, 37-45.

Tomas, F.M. & Potter, B.J. (1976b). Interaction between sites of magnesium absorption in the digestive tract of sheep. *Australian Journal of Agricultural Research* **27**, 437-446.

Tomas, F.M. & Potter, B.J. (1976c). The effect and site of action of potassium upon magnesium absorption in sheep. *Australian Journal of Agricultural Research* **27**, 873-880.

Wylie, M.J., Fontenot, J.P. & Greene, L.W. (1985). Absorption of magnesium and other macro-minerals in sheep infused with potassium in different parts of the digestive tract. *Journal of Animal Science* **61**, 1219-1229.

Table 1. *Dry matter (DM-g/d) intake and mineral concentration (g/kg DM) of the concentrate diet and lucerne hay*

Intake		Mineral concentration				
		Mg	K	Na	Ca	P
Concentrate	801	1.35	12.14	0.91	5.57	2.68
Lucerne Hay	92	2.32	57.56	0.29	9.95	2.67

Table 2. *Effect of potassium(K) and magnesium(Mg) infusion on mean urinary and faecal magnesium excretion (g/day) plasma magnesium concentration (mmol/l) and duodenal magnesium flow (g/day) during the balance period.*

Treatment	Urinary Mg (g/day)	Faecal Mg (g/day)	Plasma Mg (mmol/l)	Duodenal Mg Flow (g/day)
K1	0.61	1.34	0.94	1.65
K2	0.55	1.38	0.87	1.86
K3	0.47	1.47	0.81	2.08
K4	0.42	1.43	0.79	2.02
sed	0.022	0.094	0.023	0.110
df	12	12	12	8
<i>P</i>	<0.001	n.s.	<0.001	<0.05
Mg1	0.28	0.85	0.81	1.17
Mg2	0.42	1.18	0.84	1.52
Mg3	0.60	1.45	0.88	2.04
Mg4	0.74	2.14	0.88	2.88
sed	0.030	0.069	0.014	0.052
df	24	24	24	12
<i>P</i>	<0.001	<0.001	0.001	<0.001

Table 3: *Effect of potassium and magnesium infusion on rumen digesta pH, rumen Mg solubility and rumen whole digesta and supernatant Mg concentration.*

Treatment	Digesta pH	Rumen Mg solubility (%)	Mg concentration whole rumen digesta mg/100g DM	Mg concentration in supernatant mg/100 ml
K1	5.83	37	224	26.8
K2	5.71	35	246	27.8
K3	6.02	34	256	24.8
K4	5.91	33	265	23.7
sed	0.085	5.3	14.3	3.55
df	12	8	8	8
<i>P</i>	<0.05	n.s.	n..s.	n.s.
Mg 1	5.88	34	162	19
Mg 2	5.85	33	211	21
Mg 3	5.90	36	264	26
Mg 4	5.84	36	353	37
sed	0.044	1.4	8.1	1.2
df	24	12	12	12
<i>P</i>	n.s.	n.s.	<0.001	<0.001

Figure 1: Effect of potassium and magnesium infusions on the partitioning of magnesium absorption between the rumen and a post ruminal site.

