

Nucleotide Sequence Variation in the Insulin-Like Growth Factor 1 Gene Affects Growth and Carcass Traits in New Zealand Romney Sheep

Shaobin Li,^{1,2} Huitong Zhou,^{1–3} Fangfang Zhao,^{1,2} Qian Fang,^{2,3} Jiqing Wang,^{1,2}
Xiu Liu,^{1,2} Yuzhu Luo,^{1,2} and Jon G.H. Hickford^{2,3}

Insulin-like growth factor 1 (IGF1) is a mediator of the effects of growth hormone and polymorphism in the IGF1 gene (*IGF1*) is reported to affect fat deposition in some livestock species. In this study, nucleotide sequence variation in three regions of ovine *IGF1* (part of the 5' flanking region, the exon 3 region, and the exon 4 region) was investigated in 848 New Zealand Romney lambs using PCR-single strand conformation polymorphism (SSCP) analyses to ascertain if single nucleotide polymorphisms (SNPs) existed. Six SNPs were identified across these three regions. The effect of the sequence variation in the exon 3 and exon 4 regions on growth and carcass traits were investigated. One of the PCR-SSCP sequence variants in the exon 3 region was associated with variation in hot carcass weight, carcass fat depth at the 12th rib measured using video imaging and the percentage proportion of leg lean meat, whereas the other was associated with variation in growth rate to weaning. No associations were detected for the other gene regions analyzed. The results suggest that polymorphism in exon 3 of ovine *IGF1* has potential for use as a gene-marker for some carcass and growth traits.

Keywords: insulin-like growth factor 1 gene, sheep, polymorphism, carcass, growth

Introduction

INSULIN-LIKE GROWTH FACTOR 1 (IGF1) is encoded by the IGF1 gene (*IGF1*) (Jansen *et al.*, 1983; Hoppener *et al.*, 1985). It has “non-suppressible insulin-like activity” (Salmon and Daughaday, 1957) and is a primary mediator of the effects of growth hormone. Growth hormone is synthesized in the anterior pituitary gland and released into the blood stream. It stimulates the liver to produce IGF1, which can then fuel body growth by having growth-promoting effects on almost every cell in the body, while also regulating cellular DNA synthesis (Yakar *et al.*, 2002).

In mammals, *IGF1* is composed of six exons separated by five introns, and it spans >80 kb (Rotwein, 2012). The nucleotide sequence and length of exons 1–4 are conserved across species, whereas exons 5 and 6 are more variable. Exons 1 and 2 determine the class of the protein and encode the signal peptide for cellular localization after translation,

whereas exons 3 and 4 primarily encode the mature IGF1 peptide. This ultimately becomes the receptor-binding ligand (Rotwein, 2012).

Polymorphism of *IGF1* has been reported to affect growth and production traits in a number of livestock species. It has been reported that a single nucleotide polymorphism (SNP) in the promoter of *IGF1* affects fat deposition and carcass merit traits in hybrid Angus and Charolais beef cattle (Islam *et al.*, 2009); and in dairy cattle, SNPs in *IGF1* have been associated with growth-related traits, carcass fat, milk production, and milk fatty acid traits (Mullen *et al.*, 2011; Li *et al.*, 2016). In pigs polymorphism of *IGF1* is associated with final body weight, average daily gain and back-fat thickness (Niu *et al.*, 2013), whereas in goats an *IGF1* SNP affects growth traits (Zhang *et al.*, 2008).

There have been a number of studies investigating the effects of *IGF1* polymorphism on growth and production traits in different sheep breeds, but the results do at times

¹Gansu Key Laboratory of Herbivorous Animal Biotechnology, Faculty of Animal Science and Technology, Gansu Agricultural University, Lanzhou, China.

²International Science and Technology Cooperation Base of Meat Sheep and Meat Cattle Genetic Improvement in Northwest of China, Gansu Agricultural University, Lanzhou, China.

³Gene-Marker Laboratory, Faculty of Agricultural and Life Sciences, Lincoln University, Lincoln, New Zealand.

conflict. Some researchers have reported polymorphism in the 5'-flanking region of *IGF1* associated with growth traits in Baluchi (Tahmoorespur *et al.*, 2009), Makui (Hajihosseini *et al.*, 2013) and Makooei sheep (Negahdary *et al.*, 2013), but no associations between *IGF1* polymorphism and growth traits were detected in Indian Madras Red sheep (Ramasamy, 2018), Polish Pomeranina coarse-wool sheep (Proskura and Szewczuk, 2014), Zandi sheep (Nazari *et al.*, 2016) and Baluchi sheep (Gholibeikifard *et al.*, 2013). With Colored Polish Merino sheep, polymorphism in the 5'-flanking region of *IGF1* not only affected growth and body size, but also affects carcass and meat quality traits (Grochowska *et al.*, 2017). SNPs in *IGF1* intron 1 were found to be associated with a number of carcass traits in Santa Ines sheep, including internal carcass length, rump girth, rib yield and neck weight (Meira *et al.*, 2019).

Despite the interest in ovine *IGF1*, research has tended to focus on SNPs in the 5' flanking region and introns. Little is known about whether nucleotide sequence variation in the other regions of *IGF1* has an effect on growth and carcass traits, and whether the effect exists in common breeds in major sheep-farming countries.

In this study, we used PCR-single strand conformation polymorphism (PCR-SSCP) analyses to search for SNPs in the *IGF1* 5' flanking region, and in the exon 3 and 4 region that encode the IGF1 mature peptide in New Zealand (NZ) Romney sheep, the most popular sheep breed in NZ. The effect of the PCR-SSCP haplotypes on growth and carcass traits was subsequently investigated.

Materials and Methods

All research involving animals was carried out in accordance with the Animal Welfare Act 1999 (NZ Government) and the collection of sheep blood drops by nicking sheep ears is covered by Section 7.5 Animal Identification of the Animal Welfare (Sheep and Beef Cattle) Code of Welfare 2010, which is a code of welfare issued under the Animal Welfare Act 1999 (NZ Government).

Sheep investigated and data collection

Eight hundred forty-eight NZ Romney lambs, the progeny of 19 unrelated industry-sourced rams that were part of a progeny test on a commercial farm, were investigated. The gender, birth weight, birth rank (i.e., whether they were a single, twin, or triplet), and rearing rank were recorded for each lamb. All the lambs were weaned at ~90 days of age, weighed, and separated based on gender into two

mobs. The preweaning growth rate of the lambs was calculated as the average daily weight gain (grams/day) from birth to weaning.

As most of the female lambs were kept as ewe replacements for the larger commercial base flock, the draft weight and carcass data were only available from male lambs and a small number of cull ewe lambs. Lambs weighing >37 kg were first drafted for slaughter at around 16 weeks of age, with a second draft at ~20 weeks of age. All remaining male lambs were slaughtered at ~24 weeks of age. Draft weight and draft age were recorded for each lamb.

Hot carcass weights (HCWs) were measured directly on the processing chain (Alliance Food Limited, Smithfield, Timaru, NZ), which is the weight in kilograms of the carcass minus the head, gut, and pelt. Video image analysis (VIAScan; Sastek, Australia), developed by Meat and Livestock Australia and described by Hopkins *et al.* (2004), was used to estimate the following carcass traits: lean meat yield (expressed as a percentage of HCW) in the shoulder (shoulder yield), loin (loin yield) and leg (leg yield), and total yield (the sum of the shoulder, loin and leg yields for any given carcass), and V-GR (a VIAScan assessment of subcutaneous fat depth near the 12th rib). To describe the distribution of lean meat across the carcass, the proportion of total yield of shoulder, loin, or leg was also recorded, this being the yield of the specific part of the carcass divided by the total yield and expressed as a percentage.

At tailing, blood samples from all these sheep were collected onto TFN paper (Munktell Filter AB, Sweden) by nicking the lamb's ears and genomic DNA was then purified for PCR analysis using a two-step procedure described by Zhou *et al.* (2006).

PCR primers and amplification of ovine *IGF1*

Three pairs of PCR primers were designed manually to amplify a portion of the 5'-flanking region, the entirety of the exon 3 region (including parts of its flanking introns) and the exon 4 region (including parts of its flanking introns) of *IGF1* (Table 1). The PCR primers were chosen based on analysis of the ovine genome sequence Oar_v4.0 NC_019475.2, and checked for suitability as primers using DNAMAN (version 5.2.10; Lynnon BioSoft, Vaudreuil, Canada). The primers were synthesized by Integrated DNA Technologies (Coralville, IA).

The PCR amplifications were carried out using S1000 thermal cyclers (Bio-Rad, Hercules, CA), and were

TABLE 1. PCR PRIMERS AND PCR-SINGLE STRAND CONFORMATION POLYMORPHISM CONDITIONS FOR THE ANALYSIS OF OVINE *IGF1*

Region amplified	Primer sequence (5'-3')	Predicted amplicon size	PCR annealing temperature	SSCP condition
5' flanking	CAGTTGGCTTTACAGCTCAG CATCTGCTAATACACCTTACC	340 bp	60°C	25°C, 270 V, 15 h
Exon 3	CTGCTCAGAGGTCACCTCAC GCTGAAACACTAGGCTCGC	452 bp	62°C	31°C, 250 V, 19 h
Exon 4	GACTGCTGGAGATATACTGG CTGGTGGGCTTACCTTCTG	389 bp	62°C	28°C, 250 V, 15 h

IGF1, insulin-like growth factor 1; SSCP, single strand conformation polymorphism.

performed in a 15- μ L reaction containing the purified genomic DNA on a 1.2-mm punch of the TFN paper, 0.5 U Taq DNA polymerase (Qiagen, Hilden, Germany), 0.25 μ M of each primer, 2.5 mM Mg^{2+} , 150 μ M of each dNTP (Bioline, London, UK) and 1 \times the reaction buffer supplied with the enzyme. The thermal profile for amplification consisted of 2 min at 94°C, followed by 35 cycles of 30 s at 94°C, 30 s at the annealing temperatures shown in Table 1, and 30 s at 72°C; with a final extension of 5 min at 72°C.

Screening for sequence variation and sequencing of PCR-SSCP haplotypes

The PCR amplicons were screened for nucleotide sequence variation using SSCP analysis. A 0.7- μ L aliquot of each amplicon was mixed with 7 μ L of loading dye (98% formamide, 10 mM EDTA, 0.025% bromophenol blue, and 0.025% xylene-cyanol). After denaturation at 95°C for 5 min, the samples were cooled on wet ice and then loaded on 16 cm \times 18 cm, 14% acrylamide:bisacrylamide (37.5:1) (Bio-Rad) gels. Electrophoresis was performed using Protean II xi cells (Bio-Rad) in 0.5 \times TBE buffer, and the electrophoretic conditions shown in Table 1. Gels were silver stained according to the method of Byun *et al.* (2009).

The PCR amplicons identified as homozygous by SSCP analysis were directly sequenced at the Lincoln University Sequencing Facility, NZ. Sequence alignments, translations, and comparisons were carried out using DNAMAN. The SNPs that were revealed were named using the nomenclature described at www.hgvs.org/mutnomen/ and aligned to GenBank NC_019475.2 (*Ovis aries* breed Texel chromosome 18), Oar_v4.0.

Statistical analyses

There were some missing data, and sheep with incomplete records were removed from some analyses. Sample numbers, therefore, vary in different analyses. Statistical analyses were performed using Minitab version 17 (Minitab, Inc., State College, PA).

Two types of General Linear Mixed-Effects Models (GLMMs) were used to ascertain the effect of *IGF1* genotype on the measured traits. The first models ascertained the effect of the presence/absence (recorded as 1 and 0) of the PCR-SSCP variant sequences on the measured traits. The second models were pairwise comparisons between genotypes using a Tukey test with Bonferroni corrections. The core model for these analyses was $Y_{ijklm} = \mu + S_i + G_j + B_k + D_l + V_m + e_{ijklm}$, where Y_{ijklm} is the trait measured on each animal (birth weight, etc.), μ is the mean for the trait, S_i is the random effect of sire, G_j is the effect of gender, B_k is the effect of birth weight, birth rank, or rearing rank, D_l is draft age, V_m is the fixed effect of genotype or the presence/absence of each variant, and e_{ijklm} is the random residual error. For the birth weight GLMM, gender and birth rank were fitted into the models as fixed factors, but with the growth to weaning GLMMs, gender and rearing rank were fitted into the models. For carcass and yield traits, gender, birth weight, and draft age were fitted into the models as covariates. Only main effects were tested, and associations were considered significant at the 5% level.

Results

Nucleotide sequence variation in ovine *IGF1*

Two unique PCR-SSCP banding patterns were detected in each region of ovine *IGF1*, with either one or a combination of two banding patterns observed for each sheep (Fig. 1). DNA sequencing revealed that these PCR-SSCP patterns represented six unique sequences of *IGF1*. The six sequences have been deposited into GenBank with accession numbers MH144564–MH144569. In total, six SNPs were identified (Fig. 1). There was only one SNP in the exon 3 coding region, which was a synonymous SNP c.153T>C. The frequencies of these sequences in the NZ Romney sheep investigated are illustrated in Figure 1.

Effect of sequence variation in *IGF1* on carcass and growth traits

In the 5' flanking region one sequence (B_1 ; c.-648C and c.-646A) was predominant, whereas the other sequence (A_1 ; c.-648G and c.-646G) occurred at a frequency of <5%, hence the association analyses were only undertaken for the exon 3 and exon 4 regions.


For the exon 3 region, an effect of the presence/absence of PCR-SSCP variant was observed for growth rate to weaning (Table 2), with the presence of B_2 being associated ($p=0.048$) with a lower growth rate (present: 384.9 ± 3.77 grams/day; absent: 396.0 ± 5.44 grams/day). An effect of the presence/absence of PCR-SSCP variant was also observed for HCW, V-GR, and proportion leg yield (Table 2), with the presence of B_2 being associated with increased HCW ($p=0.015$) and increased V-GR ($p=0.003$), but decreased proportion leg yield ($p=0.012$). For this exon 3 region, an effect of genotype was observed for HCW, V-GR, and shoulder yield (Table 3). Sheep with genotype B_2B_2 (c.64-82CC and c.153CC) had lower HCW ($p=0.010$), lower V-GR ($p=0.005$) and less shoulder yield ($p=0.021$) than those sheep of genotype A_2A_2 (c.64-82AA and c.153TT) or A_2B_2 (c.64-82AC and c.153TC).

No associations were detected for the exon 4 region (results not shown).

Discussion

This is the first report describing associations between sequence variation in ovine *IGF1* and growth and carcass traits in NZ Romney lambs. There was only a single synonymous SNP detected in the coding region of *IGF1*, which is in agreement with the observation that *IGF1* is conserved among mammals and that the IGF1 protein, along with IGF2 and insulin, comprise a conserved protein family found in most mammalian species and in many other vertebrates (Rotwein, 2017). Highly conserved sequences are typically associated with proteins that underpin conserved or essential metabolic activities (Zhao *et al.*, 2018), and mice that are *IGF1*-null (created by homologous recombination), exhibit postnatal lethality, growth retardation, infertility, and profound defects in the development of their major organ systems, with this confirming the essential nature of the protein's activity (Liu *et al.*, 2000).

The effect of SNPs in the 5' flanking region cannot be reliably assessed in this study due to the minor sequence A (c.-648G and c.-646G) occurring at a low frequency (4.4%)



	<i>A</i> ₁	<i>B</i> ₁	SNP ID		<i>A</i> ₂	<i>B</i> ₂	SNP ID		<i>A</i> ₃	<i>B</i> ₃	SNP ID
c.-648	G	C	rs401028781	c.64-82	A	C	rs430457475	c.224-159	A	G	rs424410885
c.-646	G	A	rs422604851	c.153	T	C	rs159876393	c.224-47	A	C	rs413216906
Frequency	4.4%	95.6%		Frequency	47.2%	52.8%		Frequency	14.4%	85.6%	

FIG. 1. Sequence variation in ovine *IGF1*. Different sequences identified in three regions of *IGF1* using by PCR-SSCP analysis and DNA sequencing. The nucleotide substitutions in these sequences are illustrated, together with their frequencies in the 848 sheep investigated. *IGF1*, insulin-like growth factor 1; PCR-SSCP, PCR-single strand conformation polymorphism.

TABLE 2. ASSOCIATION OF *IGF1* EXON 3 SEQUENCES WITH GROWTH AND CARCASS TRAITS IN NEW ZEALAND ROMNEY SHEEP

Trait ^a	Variant	n		Mean ± SE ^b		p
		Absent	Present	Absent	Present	
Birth weight (kg)	<i>A</i> ₂	209	506	5.68 ± 0.07	5.77 ± 0.05	0.255
	<i>B</i> ₂	144	571	5.78 ± 0.08	5.74 ± 0.05	0.659
Growth rate to weaning (grams/day)	<i>A</i> ₂	209	506	390.1 ± 5.13	387.0 ± 3.68	0.532
	<i>B</i> ₂	144	571	396.0 ± 5.44	384.9 ± 3.77	0.048
HCW (kg)	<i>A</i> ₂	127	316	16.74 ± 0.23	17.21 ± 0.20	0.015
	<i>B</i> ₂	93	350	17.22 ± 0.25	17.02 ± 0.20	0.350
V-GR (mm)	<i>A</i> ₂	127	316	7.04 ± 0.32	7.84 ± 0.27	0.003
	<i>B</i> ₂	93	350	7.82 ± 0.34	7.52 ± 0.27	0.316
Shoulder yield (%)	<i>A</i> ₂	127	316	<i>17.01 ± 0.10</i>	<i>17.18 ± 0.09</i>	<i>0.052</i>
	<i>B</i> ₂	93	350	17.13 ± 0.11	17.12 ± 0.09	0.974
Loin yield (%)	<i>A</i> ₂	127	316	14.83 ± 0.10	14.92 ± 0.09	0.316
	<i>B</i> ₂	93	350	14.90 ± 0.11	14.90 ± 0.09	0.869
Leg yield (%)	<i>A</i> ₂	127	316	22.19 ± 0.14	22.13 ± 0.12	0.591
	<i>B</i> ₂	93	350	22.17 ± 0.15	22.15 ± 0.12	0.857
Total yield (%)	<i>A</i> ₂	127	316	54.03 ± 0.29	54.22 ± 0.24	0.415
	<i>B</i> ₂	93	350	54.20 ± 0.31	54.15 ± 0.24	0.874
Proportion shoulder yield (%)	<i>A</i> ₂	127	316	<i>31.49 ± 0.13</i>	<i>31.68 ± 0.11</i>	<i>0.080</i>
	<i>B</i> ₂	93	350	31.61 ± 0.14	31.63 ± 0.11	0.843
Proportion loin yield (%)	<i>A</i> ₂	127	316	27.44 ± 0.11	27.50 ± 0.09	0.505
	<i>B</i> ₂	93	350	27.49 ± 0.11	27.48 ± 0.09	0.918
Proportion leg yield (%)	<i>A</i> ₂	127	316	41.06 ± 0.12	40.81 ± 0.10	0.012
	<i>B</i> ₂	93	350	40.90 ± 0.13	40.89 ± 0.10	0.900

^aHCW—hot carcass weight; V-GR—VIA scan fat depth at the 12th rib.

^bPredicted means and standard error of those means derived from GLMMs, with various factors being included in the models for different traits as described in the Materials and Methods section. $p < 0.05$ are in bold, whereas $0.05 \leq p < 0.10$ are italicized. GLMMs, General Linear Mixed-Effects Models.

TABLE 3. ASSOCIATION OF *IGF1* EXON 3 GENOTYPES WITH GROWTH AND CARCASS TRAITS IN NEW ZEALAND ROMNEY SHEEP

Trait*	Mean \pm SE**			p
	A ₂ A ₂ (n = 144)	A ₂ B ₂ (n = 362)	B ₂ B ₂ (n = 209)	
Birth weight (kg)	5.78 \pm 0.08	5.76 \pm 0.05	5.68 \pm 0.07	0.515
Growth rate to weaning (grams/day)	395.07 \pm 5.08 (n = 93)	385.96 \pm 3.54 (n = 223)	388.76 \pm 4.55 (n = 127)	0.256
HCW (kg)	17.31 \pm 0.25^a	17.17 \pm 0.20^a	16.63 \pm 0.23^b	0.010
V-GR (mm)	7.91 \pm 0.34^a	7.67 \pm 0.28^a	6.88 \pm 0.31^b	0.005
Shoulder yield (%)	17.24 \pm 0.10^a	17.24 \pm 0.08^a	17.02 \pm 0.09^b	0.021
Loin yield (%)	14.91 \pm 0.11	14.92 \pm 0.09	14.83 \pm 0.11	0.604
Leg yield (%)	22.16 \pm 0.15	22.12 \pm 0.13	22.19 \pm 0.14	0.824
Total yield (%)	54.22 \pm 0.24	54.30 \pm 0.23	53.91 \pm 0.52	0.717
Proportion shoulder yield (%)	31.79 \pm 0.12	31.83 \pm 0.11	31.62 \pm 0.12	0.148
Proportion loin yield (%)	27.46 \pm 0.10	27.38 \pm 0.08	27.35 \pm 0.10	0.648
Proportion leg yield (%)	40.76 \pm 0.11	40.79 \pm 0.10	40.97 \pm 0.11	0.126

*HCW—hot carcass weight; V-GR—VIAScan fat depth at the 12th rib.

**Predicted means and standard error of those means derived from the GLMMs, with means that do not share a superscript letter (a or b) within rows being different at $p < 0.05$ and shown in bold.

in the NZ Romney sheep investigated. However, the sequence frequencies in this region appear to be interesting. In Iranian Zandi sheep, a medium-sized dual-purpose breed used for meat and pelt production and found in the central region of Iran, those with the nucleotide sequence variation that was also revealed in A (c.-648G and c.-646G) constituted 47% of the population (Nazari *et al.*, 2016). In Colored Polish Merino sheep, c.-648G and c.-646G are common, with a frequency of 91.6% reported (Grochowska *et al.*, 2017) and it is detected at a frequency of 19.1% in Small Tail Han sheep (primarily a meat breed in China), and was very rare or absent in Texel and Dorset sheep (both meat breeds) (He *et al.*, 2012). Whether this difference in sequence frequency is related to meat/wool/pelt production, or just reflects breed differences, awaits further investigation. However, the findings that the SNPs in this region affected wool production, with A (c.-648G and c.-646G) being associated with increased clean fleece weight in Egyptian Barki sheep (Darwish *et al.*, 2017) and that *IGF1* transgenic sheep produced more clean fleece than their nontransgenic half-sibs at yearling shearing (Damak *et al.*, 1996), suggest that *IGF1* may play a role in regulating wool growth and production.

The finding of associations between polymorphism in *IGF1* and growth traits is notable. The two SNPs in the 5' flanking region described in Ramasamy (2018), Nazari *et al.* (2016), Grochowska *et al.* (2017), and in this study, were associated with growth traits in Baluchi sheep ($n = 102$; Tahmoosrespur *et al.*, 2009), Makui sheep ($n = 100$; Hajihosseini *et al.*, 2013), and Makoei sheep (number unknown; Negahdary *et al.*, 2013). In addition, Trukhachev *et al.* (2016) found associations between the SNPs c.-5363C>T, c.-5188G>C, c.-5186G>A and c.-4088G>A, and live weight, and reported that c.-91A>C had a correlation with live weight, wither height, croup height, width and length, and other physical attributes in rams. Associations with the 5' flanking SNPs could not be tested in this study. However Proskura and Szweczek (2014) and Gholibeikifard *et al.* (2013), investigated the effect of X69473.1:g271C>T (equivalent to c.153C>T

in this study) in Pomeranian Coarse-wool sheep and Baluchi sheep, respectively, and did not find any association with growth traits. Ramasamy (2018), Nazari *et al.* (2016) and Grochowska *et al.* (2017) investigated polymorphism in the 5' flanking region of *IGF1* and also did not detect any association with growth traits in different sheep breeds.

It is unknown whether the effect of *IGF1* polymorphism on growth traits is breed dependent, but given some of the associations were typically detected with small numbers of sheep and/or there was the lack of genetic background for statistical correction, caution should be taken when interpreting these results, and further investigations may be required to confirm the findings.

The associations detected for the exon 3 PCR-SSCP variants and HCW, V-GR, and shoulder yield suggest that exon 3 nucleotide sequence variation affects selected carcass traits, although Trukhachev *et al.* (2016) revealed no associations between the synonymous substitution of c.81T>C in this exon and meat production parameters.

Given that HCW and V-GR have a moderate positive correlation ($r = 0.573$; Supplementary Table S1), the associations detected for HCW and V-GR may be due to these traits being correlated. Polymorphism in the 5' flanking region of *IGF1* affected EUROP fat class, kidney fat class, and external fatness of carcass class in Colored Polish Merino sheep (Grochowska *et al.*, 2017). Another study in Mehraban sheep describes how *IGF1* polymorphism is associated with the triglyceride and cholesterol content of blood and the authors reported a tendency for association of the *IGF1* polymorphism with dorsal fat thickness (Behzadi *et al.*, 2015). In cattle, a *SnaBI* polymorphism in the regulatory region of the *IGF1* associated with subcutaneous back fat (Curi *et al.*, 2005), and a promoter SNP in *IGF1* associated with ultrasound back fat thickness and carcass average back fat in the Angus beef (Islam *et al.*, 2009). With transgenic mice, IGFI has been shown to be involved in fat cell development (Rajkumar *et al.*, 1999). The findings of this study and others suggest that *IGF1* could be considered as a candidate gene for fat-related carcass traits.

Shoulder yield only had a weak correlation with both HCW and V-GR, suggesting that whatever effect the *IGF1* polymorphism was having, it may be different to how it might affect V-GR or HCW. The effect of *IGF1* polymorphism on meat yield has been reported for both sheep and beef cattle, with Grochowska *et al.* (2017) describing how 5' flanking region polymorphism affects fore shank weight, although they did not reveal an effect on shoulder yield in the Colored Polish Merino sheep. A promoter SNP associated with carcass lean meat yield in the Angus beef population, but not in Charolais cattle and hybrid Charolais×Angus cattle (Islam *et al.*, 2009). This suggests that different SNPs in *IGF1* may have different effects on meat yield and/or the effect may vary between breeds.

The *IGF1* gene is located on ovine chromosome 3, which to date has had at least 60 quantitative trait loci (QTL) located on it (sheep QTL database, 2019; <https://www.animalgenome.org/QTLdb/sheep/>), including markers for birth weight, body weight, muscle depth, and subcutaneous fat thickness.

The genotype associations detected for HCW, V-GR, and shoulder yield suggest that B_2 is associated with a decrease in HCW, V-GR, and shoulder yield, whereas A_2 (c.64-82A and c.153T) is associated with an increase in HCW, V-GR, and shoulder yield. As there was no difference in the marginal means for these traits between A_2A_2 (c.64-82AA and c.153TT) and A_2B_2 (c.64-82AC and c.153TC) sheep, this suggests that B_2 (c.64-82C and c.153C) has a recessive effect, whereas A_2 (c.64-82A and c.153T) has a dominant effect on these traits.

The effect of A_2 (c.64-82A and c.153T) may come about directly as a result of the two SNPs. Although the SNP in the coding region (c.153T>C) was synonymous, and would not result in an amino acid substitution, it may affect the expression or structure of the protein. It has been reported that silent mutations may affect mRNA translation rates and thus potentially change the way that protein folds (Hurst, 2011). With the intronic SNP c.64-82A>C, introns are known to carry regulatory sequences, so although they may not have a direct involvement in the regulation of transcription of highly expressed genes (Mullen *et al.*, 2011), they can affect alternative splicing mechanism and may be associated with mRNA transport or chromatin assembly (Jo and Choi, 2015).

Finally, it is quite possible that the effects observed in this research are due to the SNPs observed being linked to nucleotide sequence variation in other regions of the gene that regulate gene expression or function.

Conclusions

This study used PCR-SSCP to screen for nucleotide sequence variation in the 5' flanking region, exon 3, and exon 4 regions of ovine *IGF1*. Six previously identified SNPs were identified in 848 NZ Romney sheep. In different models, sequence variation in exon 3 of *IGF1* was associated with growth rate to weaning, HCW, V-GR, and shoulder lean meat yield and proportion leg yield. Verification of these findings will require further testing in more sheep from different flocks and breeds.

Acknowledgments

The authors thank Seung OK Byun, Yunhai Li, and Lucy Burrows for technical assistance.

Disclosure Statement

No competing financial interests exist.

Funding Information

This research was funded by Discipline Construction Fund Project of Gansu Agricultural University (GAU-XKJS-2018-030).

Supplementary Material

Supplementary Table S1

References

- Behzadi, S., Miraei, A.S., Sadeghi, M., Zamani, P., and Abdoli, R. (2015). Association of IGF-1 gene polymorphisms with carcass traits in Iranian Mehraban sheep using SSCP analysis. *Iran J Appl Anim Sci* **5**, 121–126.
- Byun, S.O., Fang, Q., Zhou, H., and Hickford, J. (2009). An effective method for silver-staining DNA in large numbers of polyacrylamide gels. *Anal Biochem* **385**, 174–175.
- Curi, R.A., De Oliveira, H., Silveira, A.C., and Lopes, C. (2005). Association between IGF-I, IGF-IR and GHRH gene polymorphisms and growth and carcass traits in beef cattle. *Livestock Prod Sci* **94**, 159–167.
- Damak, S., Su H-y, Jay, N.P., and Bullock, D.W. (1996). Improved wool production in transgenic sheep expressing insulin-like growth factor 1. *Biotechnology* **14**, 185–188.
- Darwish, H., El-Shorbagy, H., Abou-Eisha, A., El-Din, A., and Farag, I. (2017). New polymorphism in the 5' flanking region of IGF-1 gene and its association with wool traits in Egyptian Barki sheep. *J Genet Eng Biotechnol* **15**, 437–441.
- Gholibeikifard, A., Aminafshar, M., and Hosseinpour, M.M. (2013). Polymorphism of IGF-I and ADRB3 genes and their association with growth traits in the Iranian Baluchi sheep. *J Agricult Sci Technol* **15**, 1153–1162.
- Grochowska, E., Borys, B., Janiszewski, P., Knapik, J., and Mroczkowski, S. (2017). Effect of the IGF-I gene polymorphism on growth, body size, carcass and meat quality traits in Coloured Polish Merino sheep. *Archiv fuer Tierzucht* **60**, 161.
- Hajihosseino, A., Hashemi, A., Razavi-Sheshdeh, S., and Pirany, N. (2013). Association of the polymorphism in the 5' flanking region of the ovine IGF-I gene with growth and development traits in Makui sheep of Iran. *Eur J Zoologic Res* **2**, 19–24.
- He, J., Zhang, B., Chu, M., Wang, P., Feng, T., Cao, G., *et al.* (2012). Polymorphism of insulin-like growth factor 1 gene and its association with litter size in Small Tail Han sheep. *Mol Biol Rep* **39**, 9801–9807.
- Hopkins, D., Safari, E., Thompson, J., and Smith, C. (2004). Video image analysis in the Australian meat industry—precision and accuracy of predicting lean meat yield in lamb carcasses. *Meat Sci* **67**, 269–274.
- Hoppener, J.W., de Pagter-Holthuizen, P., Geurts van Kessel, A.H., Jansen, M., Kittur, S.D., Antonarakis, S.E., *et al.* (1985). The human gene encoding insulin-like growth factor I is located on chromosome 12. *Hum Genet* **69**, 157–160.
- Hurst, L.D. (2011). The sound of silence. *Nature* **471**, 582–583.

- Islam, K.K., Vinsky, M., Crews, R.E., Okine, E., Moore, S.S., Crews, D.H., Jr., *et al.* (2009). Association analyses of a SNP in the promoter of IGF1 with fat deposition and carcass merit traits in hybrid, Angus and Charolais beef cattle. *Anim Genet* **40**, 766–769.
- Jansen, M., van Schaik, F.M., Ricker, A.T., Bullock, B., Woods, D.E., Gabbay, K.H., *et al.* (1983). Sequence of cDNA encoding human insulin-like growth factor I precursor. *Nature* **306**, 609–611.
- Jo, B.-S., and Choi, S.S. (2015). Introns: the functional benefits of introns in genomes. *Genom Informat* **13**, 112.
- Li, C., Sun, D., Zhang, S., Yang, S., Alim, M.A., Zhang, Q., *et al.* (2016). Genetic effects of FASN, PPARGC1A, ABCG2 and IGF1 revealing the association with milk fatty acids in a Chinese Holstein cattle population based on a post genome-wide association study. *BMC Genet* **17**, 110.
- Liu, J.-L., Yakar, S., and LeRoith, D. (2000). Conditional knockout of mouse insulin-like growth factor-1 gene using the Cre/loxP system (44500). *Proc Soc Exp Biol Med* **223**, 344–351.
- Meira, A., Montenegro, H., Coutinho, L., Mourão G, Azevedo, H., Muniz, E., *et al.* (2019). Single nucleotide polymorphisms in the growth hormone and IGF type-1 (IGF1) genes associated with carcass traits in Santa Ines sheep. *Animal* **13**, 460–468.
- Mullen, M.P., Berry, D.P., Howard, D.J., Diskin, M.G., Lynch, C.O., Giblin, L., *et al.* (2011). Single nucleotide polymorphisms in the insulin-like growth factor 1 (IGF-1) gene are associated with performance in Holstein-Friesian dairy cattle. *Front Genet* **2**, 3.
- Nazari, F., Noshary, A., and Hemati, B. (2016). Association between Insulin-like growth factor I polymorphism and early growth traits in Iranian Zandi sheep, found polymerase chain reaction restriction fragment length polymorphism (PCR-RFLP). *Iran J Appl Anim Sci* **6**, 665–669.
- Negahdary, M., Hajihosseino, A., and Ajdary, M. (2013). PCR-SSCP variation of IGF1 and PIT1 genes and their association with estimated breeding values of growth traits in Makooei sheep. *Genet Res Int* **2013**, 272346.
- Niu, P., Kim, S.W., Choi, B.H., Kim, T.H., Kim, J.J., and Kim, K.S. (2013). Porcine insulin-like growth factor 1 (IGF1) gene polymorphisms are associated with body size variation. *Genes Genom* **35**, 523–528.
- Proskura, W.S., and Szewczuk, M. (2014). The polymorphism in the IGF1R gene is associated with body weight and average daily weight gain in Pomeranian Coarsewool ewes. *Pak Vet J* **34**, 514–517.
- Rajkumar, K., Modric, T., and Murphy, L. (1999). Impaired adipogenesis in insulin-like growth factor binding protein-1 transgenic mice. *J Endocrinol* **162**, 457.
- Ramasamy, C. (2018). Association of IGF1 gene polymorphism with growth rates in Madras Red sheep. *Int J Livestock Res* **8**, 131–137.
- Rotwein, P. (2012). Mapping the growth hormone—Stat5b—IGF-I transcriptional circuit. *Trends Endocrinol Metabol* **23**, 186–193.
- Rotwein, P. (2017). Diversification of the insulin-like growth factor 1 gene in mammals. *PLoS One* **12**, e0189642.
- Salmon, W.D., Jr., and Daughaday, W.H. (1957). A hormonally controlled serum factor which stimulates sulfate incorporation by cartilage in vitro. *J Lab Clin Med* **49**, 825–836.
- Tahmoorespur, M., Valeh, M., Nassiry, M., Moussavi, A., and Ansary, M. (2009). Association of the polymorphism in the 5' flanking region of the ovine IGF-I gene with growth traits in the Baluchi sheep. *South Afr J Anim Sci* **39**, 97–101.
- Trukhachev, V., Skripkin, V., Kvochko, A., Kulichenko, A., Kovalev, D., Pisarenko, S., *et al.* (2016). Polymorphisms of the IGF1 gene in Russian Sheep breeds and their influence on some meat production parameters. *Slovenian Vet Res* **53**, 77–83.
- Yakar, S., Rosen, C.J., Beamer, W.G., Ackert-Bicknell, C.L., Wu, Y., Liu, J.L., *et al.* (2002). Circulating levels of IGF-1 directly regulate bone growth and density. *J Clin Invest* **110**, 771–781.
- Zhang, C., Zhang, W., Luo, H., Yue, W., Gao, M., and Jia, Z. (2008). A new single nucleotide polymorphism in the IGF-I gene and its association with growth traits in the Nanjiang Huang goat. *Asian Australas J Anim Sci* **21**, 1073–1079.
- Zhao, F., Zhou, H., Li, S., Fang, Q., Luo, Y., and Hickford, J.G. (2018). Growth and carcass trait association with variation in the somatostatin receptor 1 (SSTR1) gene in New Zealand Romney sheep. *N Z J Agricult Res* **61**, 477–486.
- Zhou, H., Hickford, J., and Fang, Q. (2006). A two-step procedure for extracting genomic DNA from dried blood spots on filter paper for polymerase chain reaction amplification. *Anal Biochem* **1**, 159–161.

Address correspondence to:

Yuzhu Luo, PhD
Gansu Key Laboratory of Herbivorous
Animal Biotechnology
Faculty of Animal Science and Technology
Gansu Agricultural University
Lanzhou 730070
China

E-mail: luoyz@gsau.edu.cn

Jon G.H. Hickford, PhD
Gene-Marker Laboratory
Faculty of Agricultural and Life Sciences
Lincoln University
Lincoln 7647
New Zealand

E-mail: jonathan.hickford@lincoln.ac.nz

Received for publication September 18, 2020; received in revised form November 26, 2020; accepted November 26, 2020.